

MTHSC 206 SECTION 12.4 –CROSS PRODUCTS

Kevin James

DEFINITION

Suppose that $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_2)$ are vectors. We define the cross products $a \times b$ of a and b as

$$a \times b = \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$
$$=$$

DEFINITION

Suppose that $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_3)$ are vectors. We define the cross products $a \times b$ of a and b as

$$\begin{aligned} a \times b &= \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \\ &= (a_2b_3 - a_3b_2)i + (a_3b_1 - a_1b_3)j + (a_1b_2 - a_2b_1)k. \end{aligned}$$

DEFINITION

Suppose that $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_3)$ are vectors. We define the cross products $a \times b$ of a and b as

$$\begin{aligned} a \times b &= \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \\ &= (a_2b_3 - a_3b_2)i + (a_3b_1 - a_1b_3)j + (a_1b_2 - a_2b_1)k. \end{aligned}$$

NOTE

This definition will only work for 3 dimensional vectors.

EXERCISE

① Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

PROOF.

$$(a \times b) \cdot a$$

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

PROOF.

$$\begin{aligned}(a \times b) \cdot a &= ((a_2b_3 - a_3b_2), (a_3b_1 - a_1b_3), (a_1b_2 - a_2b_1)) \cdot (a_1, a_2, a_3)\end{aligned}$$

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

PROOF.

$$\begin{aligned}(a \times b) \cdot a &= ((a_2b_3 - a_3b_2), (a_3b_1 - a_1b_3), (a_1b_2 - a_2b_1)) \cdot (a_1, a_2, a_3) \\ &= (a_2b_3 - a_3b_2)a_1 + (a_3b_1 - a_1b_3)a_2 + (a_1b_2 - a_2b_1)a_3 =\end{aligned}$$

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

PROOF.

$$\begin{aligned}(a \times b) \cdot a &= ((a_2b_3 - a_3b_2), (a_3b_1 - a_1b_3), (a_1b_2 - a_2b_1)) \cdot (a_1, a_2, a_3) \\ &= (a_2b_3 - a_3b_2)a_1 + (a_3b_1 - a_1b_3)a_2 + (a_1b_2 - a_2b_1)a_3 = 0.\end{aligned}$$

EXERCISE

- 1 Let $a = (1, 3, 2)$ and $b = (2, -1, 1)$. Compute $a \times b$.
- 2 Compute $(a \times b) \cdot a$ and $(a \times b) \cdot b$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$. Then, the vector $a \times b$ is orthogonal to a and to b .

PROOF.

$$\begin{aligned}(a \times b) \cdot a &= ((a_2b_3 - a_3b_2), (a_3b_1 - a_1b_3), (a_1b_2 - a_2b_1)) \cdot (a_1, a_2, a_3) \\ &= (a_2b_3 - a_3b_2)a_1 + (a_3b_1 - a_1b_3)a_2 + (a_1b_2 - a_2b_1)a_3 = 0.\end{aligned}$$

You can check orthogonality to b . □

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

SKETCH OF PROOF...

$$|a \times b|^2 = (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2$$

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

SKETCH OF PROOF...

$$\begin{aligned} |a \times b|^2 &= (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \\ \dots &= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2 \end{aligned}$$

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

SKETCH OF PROOF...

$$\begin{aligned} |a \times b|^2 &= (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \\ \dots &= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2 \\ &= |a|^2|b|^2 - (a \cdot b)^2 \end{aligned}$$

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

SKETCH OF PROOF...

$$\begin{aligned} |a \times b|^2 &= (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \\ \dots &= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2 \\ &= |a|^2|b|^2 - (a \cdot b)^2 \\ &= |a|^2|b|^2 - (|a||b| \cos(\theta))^2 \end{aligned}$$

THEOREM

Suppose that $a, b \in \mathbb{R}^3$ and that θ is the angle between them.
Then $|a \times b| = |a||b| \sin(\theta)$.

SKETCH OF PROOF...

$$\begin{aligned} |a \times b|^2 &= (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \\ \dots &= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2 \\ &= |a|^2|b|^2 - (a \cdot b)^2 \\ &= |a|^2|b|^2 - (|a||b| \cos(\theta))^2 \\ &= |a|^2|b|^2(1 - \cos^2(\theta)). \end{aligned}$$



COROLLARY

Two nonzero vectors $a, b \in \mathbb{R}^3$ are parallel if and only if $a \times b = 0$.

COROLLARY

Two nonzero vectors $a, b \in \mathbb{R}^3$ are parallel if and only if $a \times b = 0$.

FACT

The area of the parallelogram determined by $a, b \in \mathbb{R}^3$ is given by $|a \times b|$.

COROLLARY

Two nonzero vectors $a, b \in \mathbb{R}^3$ are parallel if and only if $a \times b = 0$.

FACT

The area of the parallelogram determined by $a, b \in \mathbb{R}^3$ is given by $|a \times b|$.

EXERCISE

Find the area of the triangle with vertices $P = (1, 1, 1)$,
 $Q = (1, 2, 1)$ and $R = (2, 2, 3)$.

NOTE

$$\begin{array}{lll} i \times j = k & j \times k = i, & k \times i = j. \\ j \times i = -k & k \times j = -i, & i \times k = -j. \end{array}$$

NOTE

$$\begin{array}{lll} i \times j = k & j \times k = i, & k \times i = j. \\ j \times i = -k & k \times j = -i, & i \times k = -j. \end{array}$$

PROPERTIES OF \times

Suppose that $a, b, c \in \mathbb{R}^3$ and $d \in \mathbb{R}$. Then,

- 1 $a \times b = -b \times a$.
- 2 $(da) \times b = d(a \times b) = a \times (db)$.
- 3 $a \times (b + c) = a \times b + a \times c$.
- 4 $(a + b) \times c = a \times c + b \times c$.
- 5 $a \cdot (b \times c) = (a \times b) \cdot c$.
- 6 $a \times (b \times c) = (a \cdot c)b - (a \cdot b)c$. In particular \times is not associative.

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

PROOF.

$$V = (\text{Area of parallelogram determined by } b \text{ and } c) * (\text{height})$$

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

PROOF.

$$\begin{aligned} V &= (\text{Area of parallelogram determined by } b \text{ and } c) * (\text{height}) \\ &= (\text{height}) * |b \times c| \end{aligned}$$

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

PROOF.

$$\begin{aligned} V &= (\text{Area of parallelogram determined by } b \text{ and } c) * (\text{height}) \\ &= (\text{height}) * |b \times c| \\ &= |\text{Proj}_{b \times c}(a)| * |b \times c| \end{aligned}$$

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

PROOF.

$$\begin{aligned} V &= (\text{Area of parallelogram determined by } b \text{ and } c) * (\text{height}) \\ &= (\text{height}) * |b \times c| \\ &= |\text{Proj}_{b \times c}(a)| * |b \times c| \\ &= \left| \frac{(b \times c) \cdot a}{|b \times c|^2} (b \times c) \right| * |b \times c| \end{aligned}$$

VOLUMES AND TRIPLE PRODUCTS

THEOREM

The volume of the parallelepiped determined by $a, b, c \in \mathbb{R}^3$ is given by $V = |a \cdot (b \times c)|$.

PROOF.

$$\begin{aligned} V &= (\text{Area of parallelogram determined by } b \text{ and } c) * (\text{height}) \\ &= (\text{height}) * |b \times c| \\ &= |\text{Proj}_{b \times c}(a)| * |b \times c| \\ &= \left| \frac{(b \times c) \cdot a}{|b \times c|^2} (b \times c) \right| * |b \times c| \\ &= \frac{|(b \times c) \cdot a|}{|b \times c|^2} |b \times c| * |b \times c| \end{aligned}$$



EXERCISE

Use the vector triple product to show that the vectors $a = (1, 1, 1)$, $b = (1, 2, 1)$ and $c = (2, 1, 2)$ are coplanar.

Suppose that a force vector F acts on one end a rigid body which is fixed at its other end and is represented by the position vector r . We define the torque vector τ as

$$\tau = r \times F.$$

Suppose that a force vector F acts on one end a rigid body which is fixed at its other end and is represented by the position vector r . We define the torque vector τ as

$$\tau = r \times F.$$

The torque vector indicates the direction of rotation (using the right-hand rule) and measures the tendency of the object to rotate about the origin.

Suppose that a force vector F acts on one end a rigid body which is fixed at its other end and is represented by the position vector r . We define the torque vector τ as

$$\tau = r \times F.$$

The torque vector indicates the direction of rotation (using the right-hand rule) and measures the tendency of the object to rotate about the origin.

Note that

$$|\tau| = |r \times F| = |r||F| \sin(\theta),$$

where θ is the angle between F and r .