

# MTHSC 206 SECTION 16.2 – LINE INTEGRALS

Kevin James

## LINE INTEGRALS

We wish to develop the notion of integrating a real valued function  $f(x, y)$  along a smooth curve.

## LINE INTEGRALS

We wish to develop the notion of integrating a real valued function  $f(x, y)$  along a smooth curve.

Suppose that  $C$  is a smooth curve which is parametrized by  $r(t) = [x(t), y(t)]$  as  $a \leq t \leq b$ .

## LINE INTEGRALS

We wish to develop the notion of integrating a real valued function  $f(x, y)$  along a smooth curve.

Suppose that  $C$  is a smooth curve which is parametrized by

$$r(t) = [x(t), y(t)] \text{ as } a \leq t \leq b.$$

For  $n \geq 1$ , we define  $\Delta t = \frac{b-a}{n}$ ,  $t_i = a + i\Delta t$  and

$$s_i = r(t_i) = [x(t_i), y(t_i)] = [x_i, y_i].$$

## LINE INTEGRALS

We wish to develop the notion of integrating a real valued function  $f(x, y)$  along a smooth curve.

Suppose that  $C$  is a smooth curve which is parametrized by  $r(t) = [x(t), y(t)]$  as  $a \leq t \leq b$ .

For  $n \geq 1$ , we define  $\Delta t = \frac{b-a}{n}$ ,  $t_i = a + i\Delta t$  and  $s_i = r(t_i) = [x(t_i), y(t_i)] = [x_i, y_i]$ .

Also, we define  $\Delta s_i$  to be the length of the arc from  $r(t_{i-1})$  to  $r(t_i)$ .

## LINE INTEGRALS

We wish to develop the notion of integrating a real valued function  $f(x, y)$  along a smooth curve.

Suppose that  $C$  is a smooth curve which is parametrized by  $r(t) = [x(t), y(t)]$  as  $a \leq t \leq b$ .

For  $n \geq 1$ , we define  $\Delta t = \frac{b-a}{n}$ ,  $t_i = a + i\Delta t$  and  $s_i = r(t_i) = [x(t_i), y(t_i)] = [x_i, y_i]$ .

Also, we define  $\Delta s_i$  to be the length of the arc from  $r(t_{i-1})$  to  $r(t_i)$ .

## DEFINITION

We define the line integral of  $f$  along  $C$  by

$$\int_C f(x, y) \, ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}) \Delta s_i$$

## NOTE

We note that  $\Delta s_i \approx |s_i - s_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2}$ .

## NOTE

We note that  $\Delta s_i \approx |s_i - s_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2}$ .  
Since  $x_i - x_{i-1} = x(t_{i-1} + \Delta t) - x(t_{i-1}) \approx x'(t_{i-1})\Delta t$  and  
similarly for  $y_i - y_{i-1}$ ,



## NOTE

We note that  $\Delta s_i \approx |s_i - s_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2}$ .  
Since  $x_i - x_{i-1} = x(t_{i-1} + \Delta t) - x(t_{i-1}) \approx x'(t_{i-1})\Delta t$  and  
similarly for  $y_i - y_{i-1}$ ,  $\Delta s_i \approx \left( \sqrt{x'(t_{i-1})^2 + y'(t_{i-1})^2} \right) \Delta t$ .

## NOTE

We note that  $\Delta s_i \approx |s_i - s_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2}$ .

Since  $x_i - x_{i-1} = x(t_{i-1} + \Delta t) - x(t_{i-1}) \approx x'(t_{i-1})\Delta t$  and

similarly for  $y_i - y_{i-1}$ ,  $\Delta s_i \approx \left( \sqrt{x'(t_{i-1})^2 + y'(t_{i-1})^2} \right) \Delta t$ .

Thus, one can prove

$$\int_C f(x, y) \, ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}) \left( \sqrt{x'(t_{i-1})^2 + y'(t_{i-1})^2} \right) \Delta t$$

## NOTE

We note that  $\Delta s_i \approx |s_i - s_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2}$ .  
Since  $x_i - x_{i-1} = x(t_{i-1} + \Delta t) - x(t_{i-1}) \approx x'(t_{i-1})\Delta t$  and  
similarly for  $y_i - y_{i-1}$ ,  $\Delta s_i \approx \left( \sqrt{x'(t_{i-1})^2 + y'(t_{i-1})^2} \right) \Delta t$ .  
Thus, one can prove

$$\int_C f(x, y) \, ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}) \left( \sqrt{x'(t_{i-1})^2 + y'(t_{i-1})^2} \right) \Delta t$$

## FACT

$$\int_C f(x, y) \, ds = \int_a^b f(x(t), y(t)) \sqrt{(x'(t))^2 + (y'(t))^2} \, dt.$$

### EXAMPLE

Evaluate  $\int_C (2 + x^2y) \, ds$  where  $C$  is the upper half of the unit circle.

### EXAMPLE

Evaluate  $\int_C (2 + x^2y) \, ds$  where  $C$  is the upper half of the unit circle.

### EXAMPLE

Evaluate  $\int_C (x^2 + y^2) \, ds$  along the straight line from  $(a, 0)$  to  $(b, 0)$ .

## DEFINITION

If  $C$  is a piecewise smooth curve, that is it is the union of smooth curves  $C_1, C_2, \dots, C_n$  where the initial point of  $C_i$  is the end point of  $C_{i-1}$ , then we define

$$\int_C f(x, y) \, ds = \sum_{i=1}^n \int_{C_i} f(x, y) \, ds.$$

## DEFINITION

If  $C$  is a piecewise smooth curve, that is it is the union of smooth curves  $C_1, C_2, \dots, C_n$  where the initial point of  $C_i$  is the end point of  $C_{i-1}$ , then we define

$$\int_C f(x, y) \, ds = \sum_{i=1}^n \int_{C_i} f(x, y) \, ds.$$

## EXAMPLE

Integrate  $f(x, y) = 2x$  along the piecewise smooth curve  $C$  given by first traveling from  $(0, 0)$  to  $(1, 1)$  along the parabola  $y = x^2$  and then traveling along the line segment from  $(1, 1)$  to  $(1, 2)$ .

## DEFINITION

If  $C$  is a piecewise smooth curve, that is it is the union of smooth curves  $C_1, C_2, \dots, C_n$  where the initial point of  $C_i$  is the end point of  $C_{i-1}$ , then we define

$$\int_C f(x, y) \, ds = \sum_{i=1}^n \int_{C_i} f(x, y) \, ds.$$

## EXAMPLE

Integrate  $f(x, y) = 2x$  along the piecewise smooth curve  $C$  given by first traveling from  $(0, 0)$  to  $(1, 1)$  along the parabola  $y = x^2$  and then traveling along the line segment from  $(1, 1)$  to  $(1, 2)$ .

## EXAMPLE

Suppose that a wire lies along the upper half of the unit circle and has linear density at any point  $(x, y)$  proportional to its distance from the line  $y = 1$ . Compute the mass and center of mass of the wire.



## DEFINITION

Two other line integrals which occur naturally in many settings are the line integrals of  $f(x, y)$  with respect to  $x$  and the line integral of  $f(x, y)$  with respect to  $y$ . These are defined as

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t))x'(t) dt \quad \text{and}$$
$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t))y'(t) dt.$$

## DEFINITION

Two other line integrals which occur naturally in many settings are the line integrals of  $f(x, y)$  with respect to  $x$  and the line integral of  $f(x, y)$  with respect to  $y$ . These are defined as

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t))x'(t) dt \quad \text{and}$$
$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t))y'(t) dt.$$

## NOTATION

Since the line integrals of functions with respect to  $x$  and  $y$  frequently appear together, we often write

$$\int_C P(x, y) dx + \int_C Q(x, y) dy = \int_C P(x, y) dx + Q(x, y) dy.$$

## EXAMPLE

Evaluate  $\int_C y^2 dx + x dy$  first where  $C$  is the line segment from  $(-5, -3)$  to  $(0, 2)$  and again where  $C$  is the arc from  $(-5, -3)$  to  $(0, 2)$  along the parabola  $x = 4 - y^2$ . Are the answers the same?

### EXAMPLE

Evaluate  $\int_C y^2 dx + x dy$  first where  $C$  is the line segment from  $(-5, -3)$  to  $(0, 2)$  and again where  $C$  is the arc from  $(-5, -3)$  to  $(0, 2)$  along the parabola  $x = 4 - y^2$ . Are the answers the same?

### NOTE

If  $C$  denotes a curve segment parametrized by  $r(t)$ ,  $a \leq t \leq b$ , then we denote by  $-C$  the same curve with opposite orientation, that is  $-C$  is parametrized by  $s(t) = r((b - t) + a)$ .

## FACT

$$\int_{-C} f(x, y) dx = - \int_C f(x, y) dx,$$
$$\int_{-C} f(x, y) dy = - \int_C f(x, y) dy$$
$$\int_{-C} f(x, y) ds = \int_C f(x, y) ds.$$

*This is because  $\Delta x$  and  $\Delta y$  can be negative when orientation is reversed as in single variable calculus. However,  $\Delta s$  was defined to be an arc length which is positive. Thus the line integral with respect to arc length is independent to orientation.*

## DEFINITION

If  $f(x, y, z)$  is a function of 3 variables and  $C$  is a smooth curve in  $\mathbb{R}^3$  parametrized by  $r(t) = [x(t), y(t), z(t)]$  for  $a \leq t \leq b$ , then we define the line integral of  $f$  with respect to arc length as

$$\int_C f(x, y, z) \, ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}, z_{i-1}) \Delta s_i$$

## DEFINITION

If  $f(x, y, z)$  is a function of 3 variables and  $C$  is a smooth curve in  $\mathbb{R}^3$  parametrized by  $r(t) = [x(t), y(t), z(t)]$  for  $a \leq t \leq b$ , then we define the line integral of  $f$  with respect to arc length as

$$\begin{aligned}\int_C f(x, y, z) \, ds &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}, z_{i-1}) \Delta s_i \\ &= \int_a^b f(x(t), y(t), z(t)) \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} \, dt\end{aligned}$$

## DEFINITION

If  $f(x, y, z)$  is a function of 3 variables and  $C$  is a smooth curve in  $\mathbb{R}^3$  parametrized by  $r(t) = [x(t), y(t), z(t)]$  for  $a \leq t \leq b$ , then we define the line integral of  $f$  with respect to arc length as

$$\begin{aligned}\int_C f(x, y, z) \, ds &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}, z_{i-1}) \Delta s_i \\ &= \int_a^b f(x(t), y(t), z(t)) \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} \, dt \\ &= \int_a^b f(r(t)) |r'(t)| \, dt.\end{aligned}$$



## DEFINITION

If  $f(x, y, z)$  is a function of 3 variables and  $C$  is a smooth curve in  $\mathbb{R}^3$  parametrized by  $r(t) = [x(t), y(t), z(t)]$  for  $a \leq t \leq b$ , then we define the line integral of  $f$  with respect to arc length as

$$\begin{aligned} \int_C f(x, y, z) \, ds &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1}, y_{i-1}, z_{i-1}) \Delta s_i \\ &= \int_a^b f(x(t), y(t), z(t)) \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} \, dt \\ &= \int_a^b f(r(t)) |r'(t)| \, dt. \end{aligned}$$

## EXAMPLE

Evaluate  $\int_C y \sin(z) \, ds$  where  $C$  is the curve parametrized by  $r(t) = [\cos(t), \sin(t), t]$  for  $0 \leq t \leq 2\pi$ .

# LINE INTEGRALS OF VECTOR FIELDS

## NOTE

Suppose that  $F$  is a continuous force field and that a particle is moved by  $F$  through a smooth curve  $C$  parametrized by  $r(t)$ ,  $a \leq t \leq b$ . Then the work done by  $F$  is given by

$$W = \int_C F \cdot T \, ds,$$

where  $T(t)$  is the unit tangent vector of  $C$  at  $r(t)$ .

## DEFINITION

Let  $F$  be a continuous vector field defined on a smooth curve  $C$  parametrized by  $r(t)$  for  $a \leq t \leq b$ . Let  $T(t)$  denote the unit tangent vector of  $r(t)$ . Then the line integral of  $F$  along  $C$  is

$$\int_C F \cdot dr = \int_a^b F(r(t)) \cdot r'(t) dt = \int_C F \cdot T ds.$$

## DEFINITION

Let  $F$  be a continuous vector field defined on a smooth curve  $C$  parametrized by  $r(t)$  for  $a \leq t \leq b$ . Let  $T(t)$  denote the unit tangent vector of  $r(t)$ . Then the line integral of  $F$  along  $C$  is

$$\int_C F \cdot dr = \int_a^b F(r(t)) \cdot r'(t) dt = \int_C F \cdot T ds.$$

## EXAMPLE

Evaluate  $\int_C F \cdot dr$  where  $F(x, y, z) = [xy, yz, zx]$  and  $C$  is parametrized by  $r(t) = [t, t^2, t^3]$  for  $0 \leq t \leq 1$ .

## NOTE

Suppose that  $F(x, y, z) = Pi + Qj + Rk$  is a vector field and that  $C$  is parametrized by  $r(t) = [x(t), y(t), z(t)]$ . Then

$$\int_C F \cdot dr = \int_C P dx + Q dy + R dz.$$