

MTHSC 206 SECTION 13.5 –EQUATIONS OF LINES AND PLANES

Kevin James

DEFINITION

A line in \mathbb{R}^3 can be described by a point and a direction vector. Given the point r_0 and the direction vector v . Any point r on the line through r_0 and parallel to v satisfies $r = r_0 + tv$ for some $t \in \mathbb{R}$.

DEFINITION

A line in \mathbb{R}^3 can be described by a point and a direction vector. Given the point r_0 and the direction vector v . Any point r on the line through r_0 and parallel to v satisfies $r = r_0 + tv$ for some $t \in \mathbb{R}$.

DEFINITION

If we let $r_0 = (x_0, y_0, z_0)$ and $v = (a, b, c)$, then the line described above could also be described by the parametric equations

$$x = x_0 + at, \quad y = y_0 + bt \quad \text{and} \quad z = z_0 + tc.$$

DEFINITION

If the coordinates of the direction vector are all non zero, then we have a third description of the line given by solving each of the parametric equations for t and equating, namely

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

DEFINITION

If the coordinates of the direction vector are all non zero, then we have a third description of the line given by solving each of the parametric equations for t and equating, namely

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

EXERCISE

Give vector, parametric and symmetric equations for the line passing through the points $A = (1, 1, 1)$ and $B = (0, -1, 2)$. At what point does this line intersect the xy -plane?

DEFINITION

A plane in \mathbb{R}^3 can be described by a point r_0 in the plane along with a normal vector n which is orthogonal to all vectors in the plane. Given this information, we see that r is in the plane if and only if $r - r_0$ is orthogonal to n , that is

$$\begin{aligned}n \cdot (r - r_0) &= 0, \quad \text{or} \\n \cdot r &= n \cdot r_0.\end{aligned}$$

Either of these equations is called the vector equation of the plane.

DEFINITION

A plane in \mathbb{R}^3 can be described by a point r_0 in the plane along with a normal vector n which is orthogonal to all vectors in the plane. Given this information, we see that r is in the plane if and only if $r - r_0$ is orthogonal to n , that is

$$\begin{aligned}n \cdot (r - r_0) &= 0, \quad \text{or} \\n \cdot r &= n \cdot r_0.\end{aligned}$$

Either of these equations is called the vector equation of the plane.

NOTE

Letting $n = (a, b, c)$, $r = (x, y, z)$ and $r_0 = (x_0, y_0, z_0)$, we obtain the scalar equation of the plane through the point $r_0 = (x_0, y_0, z_0)$ with normal vector $n = (a, b, c)$, namely

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

NOTE

The last equation can be rewritten in the form

$$ax + by + cz + d = 0.$$

This is called a linear equation of the plane.

NOTE

The last equation can be rewritten in the form

$$ax + by + cz + d = 0.$$

This is called a linear equation of the plane.

EXERCISE

Find an equation for the plane containing the points $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$.

FACT

- ① *Two planes are parallel if and only if their normal vectors are parallel.*
- ② *The angle between two intersecting planes is equal to the angle between their normal vectors.*

FACT

- 1 *Two planes are parallel if and only if their normal vectors are parallel.*
- 2 *The angle between two intersecting planes is equal to the angle between their normal vectors.*

EXERCISE

Find the angle between the planes $x + y + z = 0$ and $x + 2y + 3z = 0$. Describe their intersection.

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$\vec{b} = \vec{P_0P} =$$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$\vec{b} = \vec{P_0P} = (x_1 - x_0, y_1 - y_0, z_1 - z_0).$$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$b = \vec{P_0P} = (x_1 - x_0, y_1 - y_0, z_1 - z_0).$$

Then, $D = |\text{comp}_n(b)| =$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$b = \vec{P_0P} = (x_1 - x_0, y_1 - y_0, z_1 - z_0).$$

$$\text{Then, } D = |\text{comp}_n(b)| = \frac{|n \cdot b|}{|n|} =$$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$b = \vec{P_0P} = (x_1 - x_0, y_1 - y_0, z_1 - z_0).$$

$$\text{Then, } D = |\text{comp}_n(b)| = \frac{|n \cdot b|}{|n|} = \frac{|a(x_1 - x_0) + b(y_1 - y_0) + c(z_1 - z_0)|}{\sqrt{a^2 + b^2 + c^2}}.$$

FACT

The distance from the point $P = (x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

PROOF.

Note that a normal vector to the plane is $n = (a, b, c)$.

Let P_0 be any point in the plane and let

$$b = \vec{P_0P} = (x_1 - x_0, y_1 - y_0, z_1 - z_0).$$

$$\text{Then, } D = |\text{comp}_n(b)| = \frac{|n \cdot b|}{|n|} = \frac{|a(x_1 - x_0) + b(y_1 - y_0) + c(z_1 - z_0)|}{\sqrt{a^2 + b^2 + c^2}}.$$

Now note that $-ax_0 - by_0 - cz_0 = d$, since P_0 is in the plane. \square