

MTHSC 206 SECTION 15.5 – THE CHAIN RULE

Kevin James

THEOREM (CHAIN RULE - CASE 1)

Suppose that $z = f(x, y)$ is a differentiable function and that $x(t)$ and $y(t)$ are both differentiable functions as well. Then,

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

THEOREM (CHAIN RULE - CASE 1)

Suppose that $z = f(x, y)$ is a differentiable function and that $x(t)$ and $y(t)$ are both differentiable functions as well. Then,

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

EXAMPLE

Suppose that $z = xy^2 + 5x^3y$ where $x(t) = e^t$ and $y(t) = \sin(t)$. Find $\frac{dz}{dt}$ when $t = 0$.

THEOREM (CHAIN RULE - CASE 1)

Suppose that $z = f(x, y)$ is a differentiable function and that $x(t)$ and $y(t)$ are both differentiable functions as well. Then,

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

EXAMPLE

Suppose that $z = xy^2 + 5x^3y$ where $x(t) = e^t$ and $y(t) = \sin(t)$. Find $\frac{dz}{dt}$ when $t = 0$.

EXAMPLE

The pressure (in kilopascals kPa), volume (in liters L) and temperature (in kelvins K) of an ideal gas are related by the equation $PV = 8.31T$. Find the rate at which the pressure is changing when the temperature is 300 K and increasing at a rate of 0.1 K/s and the volume is 100 L and increasing at 0.2 L/s.

THEOREM (CHAIN RULE - CASE 2)

Suppose that $z = f(x, y)$ is a differentiable function of x and y , where $x(s, t)$ and $y(s, t)$ are also differentiable functions. Then

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s} \quad \text{and} \quad \frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

THEOREM (CHAIN RULE - CASE 2)

Suppose that $z = f(x, y)$ is a differentiable function of x and y , where $x(s, t)$ and $y(s, t)$ are also differentiable functions. Then

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s} \quad \text{and} \quad \frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

EXAMPLE

Suppose that $z = \cos(x) \sin(y)$ and $x(s, t) = st$ and $y(s, t) = s^2 t$. Compute the partial derivatives of z with respect to s and t .

THEOREM (CHAIN RULE - GENERAL VERSION)

Suppose that u is a differentiable function in the variables x_1, x_2, \dots, x_n and each x_i is a differentiable function of the variables t_1, t_2, \dots, t_m . Then,

$$\frac{\partial u}{\partial t_i} = \sum_{j=1}^n \frac{\partial u}{\partial x_j} \frac{\partial x_j}{\partial t_i}.$$

THEOREM (CHAIN RULE - GENERAL VERSION)

Suppose that u is a differentiable function in the variables x_1, x_2, \dots, x_n and each x_j is a differentiable function of the variables t_1, t_2, \dots, t_m . Then,

$$\frac{\partial u}{\partial t_i} = \sum_{j=1}^n \frac{\partial u}{\partial x_j} \frac{\partial x_j}{\partial t_i}.$$

EXAMPLE

Suppose that $u = x^3y + y^3z + z^3x$ where $x = rs \sin(t)$, $y = rs \cos(t)$ and $z = rse^t$. Find $\frac{\partial u}{\partial s}$.

IMPLICIT DIFFERENTIATION

Given an equation $F(x, y) = 0$, we suppose that this equation implicitly defines y as a function of x .

IMPLICIT DIFFERENTIATION

Given an equation $F(x, y) = 0$, we suppose that this equation implicitly defines y as a function of x .

Applying the chain rule gives

$$\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} = 0$$

IMPLICIT DIFFERENTIATION

Given an equation $F(x, y) = 0$, we suppose that this equation implicitly defines y as a function of x .

Applying the chain rule gives

$$\begin{aligned}\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} &= 0 \\ \Rightarrow \frac{dy}{dx} &= \frac{-\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial y}} = \frac{-F_x}{F_y}\end{aligned}$$

IMPLICIT DIFFERENTIATION

Given an equation $F(x, y) = 0$, we suppose that this equation implicitly defines y as a function of x .

Applying the chain rule gives

$$\begin{aligned}\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} &= 0 \\ \Rightarrow \frac{dy}{dx} &= \frac{-\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial y}} = \frac{-F_x}{F_y}\end{aligned}$$

EXAMPLE

Find the slope of the line tangent to the unit circle at the point $(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$.

Now suppose that z is given implicitly as a function of x and y by the equation

$$F(x, y, z) = 0$$

Now suppose that z is given implicitly as a function of x and y by the equation

$$\begin{aligned} F(x, y, z) &= 0 \\ \Rightarrow F(x, y, f(x, y)) &= 0 \end{aligned}$$

Now suppose that z is given implicitly as a function of x and y by the equation

$$\begin{aligned} F(x, y, z) &= 0 \\ \Rightarrow F(x, y, f(x, y)) &= 0 \\ \Rightarrow \frac{\partial F}{\partial x} \frac{\partial x}{\partial x} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} &= 0 \end{aligned}$$

Now suppose that z is given implicitly as a function of x and y by the equation

$$\begin{aligned} F(x, y, z) &= 0 \\ \Rightarrow F(x, y, f(x, y)) &= 0 \\ \Rightarrow \frac{\partial F}{\partial x} \frac{\partial x}{\partial x} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} &= 0 \\ \Rightarrow \frac{\partial F}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} &= 0 \end{aligned}$$

Now suppose that z is given implicitly as a function of x and y by the equation

$$\begin{aligned} F(x, y, z) &= 0 \\ \Rightarrow F(x, y, f(x, y)) &= 0 \\ \Rightarrow \frac{\partial F}{\partial x} \frac{\partial x}{\partial x} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} &= 0 \\ \Rightarrow \frac{\partial F}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} &= 0 \\ \Rightarrow \frac{\partial z}{\partial x} &= \frac{-\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}} \\ &= \frac{-F_x}{F_z} \end{aligned}$$

IMPLICIT DIFFERENTIATION

Suppose that x , y and z satisfy the equation $F(x, y, z) = 0$ where F is differentiable then under the assumption that z is implicitly defined as a differentiable function of x and y , we obtain the formulas

$$\frac{\partial z}{\partial x} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}}$$

$$\frac{\partial z}{\partial y} = -\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}}$$

IMPLICIT DIFFERENTIATION

Suppose that x , y and z satisfy the equation $F(x, y, z) = 0$ where F is differentiable then under the assumption that z is implicitly defined as a differentiable function of x and y , we obtain the formulas

$$\frac{\partial z}{\partial x} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}}$$

$$\frac{\partial z}{\partial y} = -\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}}$$

EXAMPLE

Recall that the equation of the unit sphere is given by $x^2 + y^2 + z^2 = 1$. Use implicit differentiation to find the equation of the tangent plane at the point $\left(\frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3}\right)$