

MTHSC 3110 SECTION 1.8 – LINEAR TRANSFORMATIONS

Kevin James

We are going to talk about special kinds of functions from \mathbb{R}^n to \mathbb{R}^m (note: n comes before m here!)

First: what is a function?

DEFINITION

We write $f : A \rightarrow B$ to denote that f is a function from the set A to the set B . What this means is that f is a rule, which assigns, for each element $a \in A$, a unique element $b \in B$ so that $b = f(a)$.

We are going to talk about special kinds of functions from \mathbb{R}^n to \mathbb{R}^m (note: n comes before m here!)

First: what is a function?

DEFINITION

We write $f : A \rightarrow B$ to denote that f is a function from the set A to the set B . What this means is that f is a rule, which assigns, for each element $a \in A$, a unique element $b \in B$ so that $b = f(a)$. We refer to A as the domain of f , and to B as the co-domain.

We are going to talk about special kinds of functions from \mathbb{R}^n to \mathbb{R}^m (note: n comes before m here!)

First: what is a function?

DEFINITION

We write $f : A \rightarrow B$ to denote that f is a function from the set A to the set B . What this means is that f is a rule, which assigns, for each element $a \in A$, a unique element $b \in B$ so that $b = f(a)$. We refer to A as the domain of f , and to B as the co-domain.

The set

$$\{f(a) : a \in A\}$$

of values taken by the function is called the image or range of f .

We will restrict our attention to functions which interact nicely with vector addition and scalar multiplication.

We will restrict our attention to functions which interact nicely with vector addition and scalar multiplication.

DEFINITION

A *linear transformation* from \mathbb{R}^n to \mathbb{R}^m is a function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ which satisfies the following two properties: whenever $\vec{u}, \vec{v} \in \mathbb{R}^n$ and $c \in \mathbb{R}$, then

- 1 $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$
- 2 $T(c\vec{u}) = cT(\vec{u})$.

NOTE

If A is a $m \times n$ matrix, then we can define $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $T(\vec{x}) = A\vec{x}$.

NOTE

If A is a $m \times n$ matrix, then we can define $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $T(\vec{x}) = A\vec{x}$.

We have already proved that

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}) \text{ and}$$
$$T(\alpha\vec{x}) = A(\alpha\vec{x}) = \alpha A\vec{x} = \alpha T(\vec{x}) \text{ for any } \alpha \in \mathbb{R}.$$

EXAMPLE

Let

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 3 \\ 5 & -1 \end{pmatrix}$$

Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by $T(\vec{x}) = A\vec{x}$.

① $T\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right) =$

② Find all $\vec{x} \in \mathbb{R}^2$ so that $T(\vec{x}) = \begin{pmatrix} 3 \\ 5 \\ 4 \end{pmatrix}$.

EXAMPLE

Let

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Then $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T(\vec{x}) = A\vec{x}$ is called a projection:

$$T \left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \right) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \\ \\ \end{pmatrix}$$

EXAMPLE

Consider the map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{x}) = A\vec{x}$ where $A = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$. This is an example of a shear transformation. Draw the image of the box $[0, 1] \times [0, 1]$ under T to see why.

EXAMPLE

Consider the map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{x}) = A\vec{x}$ where $A = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$. This is an example of a shear transformation. Draw the image of the box $[0, 1] \times [0, 1]$ under T to see why.

NOTE

If T is a linear transformation, then

- 1 $T(\vec{0}) = \vec{0}$.
- 2 $T(c\vec{u} + d\vec{v}) = cT(\vec{u}) + dT(\vec{v})$.
- 3 $T\left(\sum_{i=1}^p c_i \vec{v}_i\right) = \sum_{i=1}^p c_i T(\vec{v}_i)$

EXAMPLE

Define a map $T_r : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by $T\vec{x} = r\vec{x}$. If $0 < r < 1$, the map is called a *contraction*. If $r > 1$ it is called a *dilation*. Show that T_r is a linear transformation.