

MTHSC 3110 SECTION 1.9 – THE MATRIX OF A LINEAR TRANSFORMATION

Kevin James

RECALL

A function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a linear transformation if for every $\vec{u}, \vec{v} \in \mathbb{R}^n$ and $c \in \mathbb{R}$,

- 1 $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$.
- 2 $T(c\vec{u}) = cT(\vec{u})$

EXAMPLE

Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is a linear transformation, and that we know

$$T(\vec{e}_1) = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \text{and} \quad T(\vec{e}_2) = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

where $\vec{e}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\vec{e}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Give a complete description of T .

DEFINITION

Let $\vec{e}_j \in \mathbb{R}^n$ denote the vector having a 1 in the j^{th} row, and zeros elsewhere.

DEFINITION

Let $\vec{e}_j \in \mathbb{R}^n$ denote the vector having a 1 in the j^{th} row, and zeros elsewhere.

THEOREM

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then there exists a unique matrix A so that for every $\vec{x} \in \mathbb{R}^n$,

$$T(\vec{x}) = A\vec{x}.$$

In fact, A is the $m \times n$ matrix whose j^{th} column is $T(\vec{e}_j)$.
-i.e.

$$A = \left(\begin{array}{c|c|c|c} T(\vec{e}_1) & T(\vec{e}_2) & \dots & T(\vec{e}_n) \\ \hline \end{array} \right).$$

EXAMPLE

Find the matrix representing the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(\vec{x}) = 3\vec{x}$.

EXAMPLE

Find the matrix representing the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(\vec{x}) = 3\vec{x}$.

DEFINITION

- A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be onto \mathbb{R}^m if every $\vec{b} \in \mathbb{R}^m$ is the image of at least one $\vec{x} \in \mathbb{R}^n$. (T is onto provided that $\forall \vec{b} \in \mathbb{R}^m, \exists \vec{x} \in \mathbb{R}^n$ so that $T(\vec{x}) = \vec{b}$.)

EXAMPLE

Find the matrix representing the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(\vec{x}) = 3\vec{x}$.

DEFINITION

- A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be onto \mathbb{R}^m if every $\vec{b} \in \mathbb{R}^m$ is the image of at least one $\vec{x} \in \mathbb{R}^n$. (T is onto provided that $\forall \vec{b} \in \mathbb{R}^m, \exists \vec{x} \in \mathbb{R}^n$ so that $T(\vec{x}) = \vec{b}$.)
- A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be one-to-one (or 1-1) if every $\vec{b} \in \mathbb{R}^m$ is the image of at most one $\vec{x} \in \mathbb{R}^n$. (That is, if $T(\vec{x}) = T(\vec{y})$ then $\vec{x} = \vec{y}$.)

EXAMPLE

Suppose that T is the linear transformation with matrix

$$\begin{pmatrix} 1 & 2 & 5 & 6 \\ 0 & 1 & 3 & 4 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$

- 1 If $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, what are m and n ?
- 2 Is T 1-1? What do you have to check?
- 3 Is T onto? What do you have to check?

THEOREM

A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is 1-1 if and only if $T\vec{x} = \vec{0}$ has only the trivial solution.

PROOF.



THEOREM

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation and let A be its matrix.

- 1 T is onto if and only if the span of the columns of A is all of \mathbb{R}^m .
- 2 T is 1-1 if and only if the columns of A are linearly independent.

PROOF.



DETECTING 1-1 AND ONTO CONDITIONS

NOTE

As a consequence of this theorem we can check whether T is 1-1 or onto by row-reducing its matrix A .

DETECTING 1-1 AND ONTO CONDITIONS

NOTE

As a consequence of this theorem we can check whether T is 1-1 or onto by row-reducing its matrix A .
If there is a pivot in every row, then T is onto.

DETECTING 1-1 AND ONTO CONDITIONS

NOTE

As a consequence of this theorem we can check whether T is 1-1 or onto by row-reducing its matrix A .

If there is a pivot in every row, then T is onto.

if there is a pivot in every column, then T is 1-1.

DETECTING 1-1 AND ONTO CONDITIONS

NOTE

As a consequence of this theorem we can check whether T is 1-1 or onto by row-reducing its matrix A .

If there is a pivot in every row, then T is onto.

if there is a pivot in every column, then T is 1-1.

EXAMPLE

Let $T \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 3x_1 + x_2 \\ 5x_1 + 7x_2 \\ x_1 + 3x_2 \end{pmatrix}$. Is T 1-1 and/or onto?