

# MTHSC 3110 SECTION 3.1 – DETERMINANTS

Kevin James

## EXERCISE

Perform row reduction on the following matrices in order to characterize invertibility.

① 
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

## EXERCISE

Perform row reduction on the following matrices in order to characterize invertibility.

① 
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

② 
$$\begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$$

## DEFINITION

Suppose that  $A$  is a  $3 \times 3$  matrix as in the previous exercise. The determinant of  $A$  is defined by

$$\Delta(A) = aei + bfg + cdh - ceg - afh - dbi.$$

## DEFINITION

Suppose that  $A$  is a  $3 \times 3$  matrix as in the previous exercise. The determinant of  $A$  is defined by

$$\Delta(A) = aei + bfg + cdh - ceg - afh - dbi.$$

## FACT

*$A$  is invertible if and only if  $\Delta(A) \neq 0$ .*

## DEFINITION

Suppose that  $A$  is a  $3 \times 3$  matrix as in the previous exercise. The determinant of  $A$  is defined by

$$\Delta(A) = aei + bfg + cdh - ceg - afh - dbi.$$

## FACT

$A$  is invertible if and only if  $\Delta(A) \neq 0$ .

## NOTE

Suppose that  $A$  is a  $3 \times 3$  matrix as before. Then,

$$\Delta(A) = a \cdot \det \begin{pmatrix} e & f \\ h & i \end{pmatrix} - b \cdot \det \begin{pmatrix} d & g \\ f & i \end{pmatrix} + c \cdot \det \begin{pmatrix} d & e \\ g & h \end{pmatrix}.$$

## DEFINITION

We define the  $ij^{\text{th}}$  minor of an  $n \times n$  matrix  $A$  as

$$A_{ij} = \begin{pmatrix} a_{1,1} & \dots & a_{1,j-1} & a_{1,j+1} & \dots & a_{1,n} \\ a_{2,1} & \dots & a_{2,j-1} & a_{2,j+1} & \dots & a_{2,n} \\ \vdots & & \vdots & \vdots & & \vdots \\ a_{i-1,1} & \dots & a_{i-1,j-1} & a_{i-1,j+1} & \dots & a_{i-1,n} \\ a_{i+1,1} & \dots & a_{i+1,j-1} & a_{i+1,j+1} & \dots & a_{i+1,n} \\ \vdots & & \vdots & \vdots & & \vdots \\ a_{n,1} & \dots & a_{n,j-1} & a_{n,j+1} & \dots & a_{n,n} \end{pmatrix}$$

## DEFINITION

We define the  $ij^{\text{th}}$  minor of an  $n \times n$  matrix  $A$  as

$$A_{ij} = \begin{pmatrix} a_{1,1} & \cdots & a_{1,j-1} & a_{1,j+1} & \cdots & a_{1,n} \\ a_{2,1} & \cdots & a_{2,j-1} & a_{2,j+1} & \cdots & a_{2,n} \\ \vdots & & \vdots & \vdots & & \vdots \\ a_{i-1,1} & \cdots & a_{i-1,j-1} & a_{i-1,j+1} & \cdots & a_{i-1,n} \\ a_{i+1,1} & \cdots & a_{i+1,j-1} & a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & & \vdots & \vdots & & \vdots \\ a_{n,1} & \cdots & a_{n,j-1} & a_{n,j+1} & \cdots & a_{n,n} \end{pmatrix}$$

## DEFINITION

Suppose that  $A$  is an  $n \times n$  matrix. Then,

$$\det(A) = \sum_{j=1}^n (-1)^{1+j} a_{1j} \cdot \det(A_{1j}).$$



### EXAMPLE

Compute the  $\det(A)$  where  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 2 & 0 & 3 & 0 \\ 1 & 2 & 0 & 3 \\ 1 & 1 & 1 & 1 \end{pmatrix}$ .

## DEFINITION

We define the  $(i, j)^{\text{th}}$  cofactor of  $A$  to be

$$C_{ij} = (-1)^{i+j} \det(A_{ij}).$$

## DEFINITION

We define the  $(i, j)^{\text{th}}$  cofactor of  $A$  to be

$$C_{ij} = (-1)^{i+j} \det(A_{ij}).$$

## THEOREM

*Suppose that  $A$  is an  $n \times n$  matrix. Then we can compute the determinant of  $A$  by expanding by cofactors along any row or column of  $A$ . That, is,*

- 1  $\det(A) = \sum_{j=1}^n a_{ij} C_{ij}$       $i$  is fixed.
- 2  $\det(A) = \sum_{i=1}^n a_{ij} C_{ij}$       $j$  is fixed.

## EXAMPLE

Compute  $\det(A)$  where  $A = \begin{pmatrix} 1 & -1 & 0 \\ 2 & 2 & 0 \\ 3 & 3 & 1 \end{pmatrix}$ .

## EXAMPLE

Compute  $\det(A)$  where  $A = \begin{pmatrix} 1 & -1 & 0 \\ 2 & 2 & 0 \\ 3 & 3 & 1 \end{pmatrix}$ .

## THEOREM

*If  $A$  is triangular, then  $\det(A) = a_{11} \cdot a_{22} \cdot \cdots \cdot a_{nn}$ .*