

MTHSC 3110 SECTION 3.2 – PROPERTIES OF DETERMINANTS

Kevin James

THEOREM

Suppose that A and B are $n \times n$ matrices and that B is obtained from A by performing a single elementary row operation \mathcal{R} .

- 1 If \mathcal{R} is $R_i \leftrightarrow R_j$ then $\det(A) = -\det(B)$.
- 2 If \mathcal{R} is $R_i + cR_j$ then $\det(A) = \det(B)$.
- 3 If \mathcal{R} is cR_i then $\det(A) = \frac{1}{c} \det(B)$.

THEOREM

Suppose that A and B are $n \times n$ matrices and that B is obtained from A by performing a single elementary row operation \mathcal{R} .

- 1 If \mathcal{R} is $R_i \leftrightarrow R_j$ then $\det(A) = -\det(B)$.
- 2 If \mathcal{R} is $R_i + cR_j$ then $\det(A) = \det(B)$.
- 3 If \mathcal{R} is cR_i then $\det(A) = \frac{1}{c} \det(B)$.

EXAMPLE

Compute $\det(A)$ where $A = \begin{pmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{pmatrix}$.

COROLLARY

- 1 $\det(E_{R_i \leftrightarrow R_j}) = -1.$
- 2 $\det(E_{R_i + cR_j}) = 1.$
- 3 $\det(E_{cR_j}) = c.$
- 4 If $A \xrightarrow{\mathcal{R}} B$, then $\det(B) = \det(E_{\mathcal{R}}) \det(A).$

COROLLARY

- 1 $\det(E_{R_i \leftrightarrow R_j}) = -1.$
- 2 $\det(E_{R_i + cR_j}) = 1.$
- 3 $\det(E_{cR_j}) = c.$
- 4 If $A \xrightarrow{\mathcal{R}} B$, then $\det(B) = \det(E_{\mathcal{R}}) \det(A).$

NOTE

Given an $n \times n$ matrix A , we can obtain a row echelon form of A , using only row replacement and row interchange operations.

COROLLARY

Suppose that A is an $n \times n$ matrix and that U is an echelon form of A obtained through the use of row replacements and row interchanges only. Let r denote the number of row interchanges used to obtain U . Then,

$$\det(A) = (-1)^r \cdot u_{11} \cdot u_{22} \cdot \cdots \cdot u_{nn}$$

COROLLARY

Suppose that A is an $n \times n$ matrix and that U is an echelon form of A obtained through the use of row replacements and row interchanges only. Let r denote the number of row interchanges used to obtain U . Then,

$$\begin{aligned} \det(A) &= (-1)^r \cdot u_{11} \cdot u_{22} \cdots u_{nn} \\ &= \begin{cases} (-1)^r * \text{product of pivots in } U & \text{if } A \text{ is invertible,} \\ 0 & \text{if } A \text{ is singular.} \end{cases} \end{aligned}$$

COROLLARY

Suppose that A is an $n \times n$ matrix and that U is an echelon form of A obtained through the use of row replacements and row interchanges only. Let r denote the number of row interchanges used to obtain U . Then,

$$\begin{aligned}\det(A) &= (-1)^r \cdot u_{11} \cdot u_{22} \cdots u_{nn} \\ &= \begin{cases} (-1)^r * \text{product of pivots in } U & \text{if } A \text{ is invertible,} \\ 0 & \text{if } A \text{ is singular.} \end{cases}\end{aligned}$$

COROLLARY

Suppose that A is an $n \times n$ matrix. Then A is invertible if and only if $\det(A) \neq 0$.

COROLLARY

If A has two rows or two columns the same, then $\det(A) = 0$ and A is singular.

COROLLARY

If A has two rows or two columns the same, then $\det(A) = 0$ and A is singular.

PROOF.



COROLLARY

If A has two rows or two columns the same, then $\det(A) = 0$ and A is singular.

PROOF.



NOTE

When computing $\det(A)$, we can exploit the presence of many zeros in A .

COROLLARY

If A has two rows or two columns the same, then $\det(A) = 0$ and A is singular.

PROOF.



NOTE

When computing $\det(A)$, we can exploit the presence of many zeros in A .

EXERCISE

Compute $\det(A)$ where $A = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 8 & -1 & 5 \\ 1 & 2 & 3 & 4 \end{pmatrix}$.



THEOREM

Suppose that A is an $n \times n$ matrix. Then A and its transpose have the same determinant (-i.e. $\det(A^T) = \det(A)$).

THEOREM

Suppose that A is an $n \times n$ matrix. Then A and its transpose have the same determinant (-i.e. $\det(A^T) = \det(A)$).

THEOREM

Suppose that A and B are $n \times n$ matrices. Then

$$\det(AB) = \det(A) \det(B).$$

THEOREM

Suppose that A is an $n \times n$ matrix. Then A and its transpose have the same determinant (-i.e. $\det(A^T) = \det(A)$).

THEOREM

Suppose that A and B are $n \times n$ matrices. Then

$$\det(AB) = \det(A) \det(B).$$

EXAMPLE

Compute $\det(A)$, $\det(B)$ and $\det(AB)$ where $A = \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}$ and

$$B = \begin{pmatrix} 1 & 2 \\ -1 & 1 \end{pmatrix}.$$