# MTHSC 3190 Section 10 Quantifiers

Kevin James



An <u>existential statement</u> asserts the existence of an object of some type with certain specified properties. Such a statement typically is preceded with an <u>existential quantifier</u> such as "there exists" or "there is".

An <u>existential statement</u> asserts the existence of an object of some type with certain specified properties. Such a statement typically is preceded with an <u>existential quantifier</u> such as "there exists" or "there is".

#### EXAMPLE

There is an integer x that is divisible by 2.

An <u>existential statement</u> asserts the existence of an object of some type with certain specified properties. Such a statement typically is preceded with an <u>existential quantifier</u> such as "there exists" or "there is".

### EXAMPLE

There is an integer x that is divisible by 2.

#### General Form

There is  $x \in A$  such that P.

An <u>existential statement</u> asserts the existence of an object of some type with certain specified properties. Such a statement typically is preceded with an <u>existential quantifier</u> such as "there exists" or "there is".

#### EXAMPLE

There is an integer x that is divisible by 2.

#### General Form

There is  $x \in A$  such that P.

#### NOTATION

The existential quantifiers <u>there is</u> or <u>there exists</u> are sometimes replaced with the notation  $\exists$ .

### PROOF TEMPLATE FOR EXISTENCE

To prove the statement " $\exists x \in A \text{ such that } P$ ", we simply need to find an example of an element of A which satisfies P. Our proof will be constructed as follows.

- **1** Select a value for the variable x by stating "Let  $x = \underline{\hspace{1cm}}$ ."
- 2 Verify that x satisfies the claimed properties P.
- **3** Close the proof with "Therefore  $x = \underline{\hspace{1cm}}$  satisfies P as claimed".

### PROOF TEMPLATE FOR EXISTENCE

To prove the statement " $\exists x \in A \text{ such that } P$ ", we simply need to find an example of an element of A which satisfies P. Our proof will be constructed as follows.

- **1** Select a value for the variable x by stating "Let x =\_\_\_\_\_."
- 2 Verify that x satisfies the claimed properties P.
- **3** Close the proof with "Therefore  $x = \underline{\hspace{1cm}}$  satisfies P as claimed".

### EXAMPLE

Show that there is  $x \in \mathbb{Z}$  such that 2|x.

# DEFINITION (UNIVERSAL QUANTIFIERS AND STATEMENTS)

A <u>universal statement</u> asserts that a certain property is satisfied by all members of a certain set (or universe). Universal statements are typically preceded by a <u>universal quantifier</u> such as "for all", "every", "all", "each", ....

# DEFINITION (UNIVERSAL QUANTIFIERS AND STATEMENTS)

A <u>universal statement</u> asserts that a certain property is satisfied by all members of a certain set (or universe). Universal statements are typically preceded by a <u>universal quantifier</u> such as "for all", "every", "all", "each", ....

### EXAMPLE

- 1 Every integer is either even or odd.
- 2 All integers are . . . .
- 3 Each integer is . . . .
- **1** Let  $x ∈ \mathbb{Z}$ , then . . . .

# DEFINITION (UNIVERSAL QUANTIFIERS AND STATEMENTS)

A <u>universal statement</u> asserts that a certain property is satisfied by all members of a certain set (or universe). Universal statements are typically preceded by a <u>universal quantifier</u> such as "for all", "every", "all", "each", ....

### EXAMPLE

- 1 Every integer is either even or odd.
- 2 All integers are . . . .
- 3 Each integer is . . . .
- 4 Let  $x \in \mathbb{Z}$ , then . . . .

### NOTATION

The symbol  $\forall$  read "for all" is often used as to denote the universal quantifier

# GENERAL FORM

The general form of a statement with a universal quantifier is  $\forall x \in A, S(x),$ 

where S(x) is a statement about x.

# GENERAL FORM

The general form of a statement with a universal quantifier is  $\forall x \in A, S(x),$ 

where S(x) is a statement about x.

### EXAMPLE

 $\forall x \in \mathbb{Z}$ , x is even or x is odd.

### GENERAL FORM

The general form of a statement with a universal quantifier is  $\forall x \in A, S(x),$ 

where S(x) is a statement about x.

### EXAMPLE

 $\forall x \in \mathbb{Z}$ , x is even or x is odd.

### Proof template for Universal Statements

To prove the statement " $\forall x \in A$ , S(x)" we proceed as follows

- Start the proof by establishing notation for an arbitrary element of A with a statement like "Let  $x \in A$ " or "Suppose  $x \in A$ ."
- 2 Prove the statement S(x) about x using an appropriate proof template.

# NEGATION OF UNIVERSAL AND EXISTENTIAL STATEMENTS

- $\neg(\exists x \in A \text{ such that } P(x)) = \forall x \in A, \neg P(x).$
- $\neg(\forall x \in A, S(x)) = \exists x \in A \text{ such that } \neg S(x).$

#### NEGATION OF UNIVERSAL AND EXISTENTIAL STATEMENTS

- $\neg(\exists x \in A \text{ such that } P(x)) = \forall x \in A, \neg P(x).$
- $\neg(\forall x \in A, S(x)) = \exists x \in A \text{ such that } \neg S(x).$

### EXAMPLE

- Express the statement "There is no integer that is both even and odd." using quantifiers.
- Express the statement "Not all integers are prime." using quantifiers.

Consider the following statements. Are they true?

- $\exists y \in \mathbb{R} \text{ such that } \forall x \in \mathbb{R}, xy = 1.$