

# MTHSC 3190 SECTION 3.13

Kevin James

**DEFINITION**

A relation is a set of ordered pairs.

## DEFINITION

A relation is a set of ordered pairs.

## EXAMPLE

$$R = \{(0, 0), (0, 1), (1, 2), 2, 5)\}$$

## DEFINITION

A relation is a set of ordered pairs.

## EXAMPLE

$$R = \{(0, 0), (0, 1), (1, 2), 2, 5)\}$$

## NOTATION

If  $(x, y) \in R$ , we say “ $x$  is related to  $y$  by  $R$ ” and we may write  $xRy$ . If  $(x, y) \notin R$  we write  $x \not R y$ .

## EXAMPLE

The  $<$  relation can be thought of as the set

$$< = \{ \dots, (-2, -1), (-2, 0), \dots, (1, 2), \dots (3, 100), \dots \}$$

## EXAMPLE

The  $<$  relation can be thought of as the set

$$< = \{ \dots, (-2, -1), (-2, 0), \dots, (1, 2), \dots (3, 100), \dots \}$$

## DEFINITION

Let  $R$  be a relation and let  $A$  and  $B$  be sets.

- 1 We say that “ $R$  is a relation on  $A$ ” provided that  $R \subseteq A \times A$ .

## EXAMPLE

The  $<$  relation can be thought of as the set

$$< = \{ \dots, (-2, -1), (-2, 0), \dots, (1, 2), \dots (3, 100), \dots \}$$

## DEFINITION

Let  $R$  be a relation and let  $A$  and  $B$  be sets.

- ① We say that “ $R$  is a relation on  $A$ ” provided that  $R \subseteq A \times A$ .
- ② We say that “ $R$  is a relation from  $A$  to  $B$ ” provided that  $R \subseteq A \times B$ .

## EXAMPLE

Let  $A = \{1, 2, 3\}$  and  $B = \{3, 4, 5\}$  and let

$$R = \{(1, 1), (2, 2), (3, 3)\},$$

$$S = \{(1, 2), (2, 3)\},$$

$$T = \{(2, 4), (3, 5)\},$$

$$U = \{(4, 1), (3, 2), (5, 3)\},$$

$$V = \{(17, 32), (0, 5), (1, 3), (2, 5)\},$$

Then,

$R$  is a relation \_\_\_\_\_

$S$  is a relation \_\_\_\_\_

$T$  is a relation \_\_\_\_\_

$U$  is a relation \_\_\_\_\_

$V$  is a relation \_\_\_\_\_



## DEFINITION

Let  $R$  be a relation. The inverse of  $R$  denoted by  $R^{-1}$  is the relation formed by reversing the order of all pairs in  $R$ .

(-i.e.  $R^{-1} = \{(y, x) : (x, y) \in R\}$ .)

## DEFINITION

Let  $R$  be a relation. The inverse of  $R$  denoted by  $R^{-1}$  is the relation formed by reversing the order of all pairs in  $R$ .  
(-i.e.  $R^{-1} = \{(y, x) : (x, y) \in R\}$ .)

## EXAMPLE

Let  $R = \{(1, 1), (1, 2), (1, 3), (2, 4), (4, 2)\}$ . What is  $R^{-1}$ ?

## DEFINITION

Let  $R$  be a relation. The inverse of  $R$  denoted by  $R^{-1}$  is the relation formed by reversing the order of all pairs in  $R$ .

(-i.e.  $R^{-1} = \{(y, x) : (x, y) \in R\}$ .)

## EXAMPLE

Let  $R = \{(1, 1), (1, 2), (1, 3), (2, 4), (4, 2)\}$ . What is  $R^{-1}$ ?

## NOTE

If  $R$  is a relation from  $A$  to  $B$ , then  $R^{-1}$  is a relation from  $B$  to  $A$ .

## PROPOSITION

*Let  $R$  be a relation. Then  $(R^{-1})^{-1} = R$ .*

## PROPOSITION

*Let  $R$  be a relation. Then  $(R^{-1})^{-1} = R$ .*

## PROOF

## DEFINITION

Let  $R$  be a relation defined on a set  $A$ .

- 1 If  $\forall x \in A, xRx$ , then  $R$  is reflexive.

## DEFINITION

Let  $R$  be a relation defined on a set  $A$ .

- ① If  $\forall x \in A, xRx$ , then  $R$  is reflexive.
- ② If  $\forall x \in A, x \not R x$ , then  $R$  is irreflexive.

## DEFINITION

Let  $R$  be a relation defined on a set  $A$ .

- ① If  $\forall x \in A, xRx$ , then  $R$  is reflexive.
- ② If  $\forall x \in A, x \not R x$ , then  $R$  is irreflexive.
- ③ If  $\forall (x, y) \in R, (y, x) \in R$ , then  $R$  is symmetric.



## DEFINITION

Let  $R$  be a relation defined on a set  $A$ .

- ① If  $\forall x \in A, xRx$ , then  $R$  is reflexive.
- ② If  $\forall x \in A, x \not R x$ , then  $R$  is irreflexive.
- ③ If  $\forall (x, y) \in R, (y, x) \in R$ , then  $R$  is symmetric.
- ④ If  $\forall x, y \in A, (xRy \wedge yRx) \Rightarrow x = y$ , then  $R$  is antisymmetric.

## DEFINITION

Let  $R$  be a relation defined on a set  $A$ .

- ① If  $\forall x \in A, xRx$ , then  $R$  is reflexive.
- ② If  $\forall x \in A, x \not R x$ , then  $R$  is irreflexive.
- ③ If  $\forall (x, y) \in R, (y, x) \in R$ , then  $R$  is symmetric.
- ④ If  $\forall x, y \in A, (xRy \wedge yRx) \Rightarrow x = y$ , then  $R$  is antisymmetric.
- ⑤ If  $\forall x, y \in A, (xRy \wedge yRz) \Rightarrow xRz$ , then  $R$  is transitive.

## EXAMPLE

Let  $A = \{1, 2, 3, 4\}$ . For each of the following relations on  $A$ , indicate which properties the relation has and does not have.

- ①  $R = \{(1, 1), (2, 2), (3, 3), (4, 4)\}$
- ②  $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 3)\}$
- ③  $T = \{(1, 2), (1, 3), (2, 3)\}$
- ④  $U = \{(1, 1), (2, 3), (3, 1), (1, 2)\}$

## EXAMPLE

Let  $A = \{1, 2, 3, 4\}$ . For each of the following relations on  $A$ , indicate which properties the relation has and does not have.

- ①  $R = \{(1, 1), (2, 2), (3, 3), (4, 4)\}$
- ②  $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 3)\}$
- ③  $T = \{(1, 2), (1, 3), (2, 3)\}$
- ④  $U = \{(1, 1), (2, 3), (3, 1), (1, 2)\}$

## EXAMPLE

Some other relations to consider.

- ①  $<$  on  $\mathbb{Z}$ .
- ②  $\leq$  on  $\mathbb{Z}$ .
- ③  $|$  on  $\mathbb{N}$ .
- ④  $|$  on  $\mathbb{Z}$ .