

# MTHSC 412 SECTION 1.4 –BINARY OPERATIONS

Kevin James

## DEFINITION

A binary operation on a nonempty set  $A$  is a mapping  $f$  from  $A \times A$  to  $A$ . That is  $f \subseteq A \times A \times A$  and  $f$  has the property that for each  $(a, b) \in A \times A$ , there is precisely one  $c \in A$  such that  $(a, b, c) \in f$ .

## NOTATION

If  $f$  is a binary operation on  $A$  and if  $(a, b, c) \in f$  then we have already seen the notation  $f(a, b) = c$ . For binary operations, it is customary to write instead

$$a f b = c,$$

or perhaps

$$a * b = c.$$

## EXAMPLE

Some binary operations on  $\mathbb{Z}$  are

①  $x * y = x + y$

②  $x * y = x - y$

③  $x * y = xy$

④  $x * y = x + 2y + 3$

⑤  $x * y = 1 + xy$

# COMMUTATIVITY AND ASSOCIATIVITY

## DEFINITION

Suppose that  $*$  is a binary operation of a nonempty set  $A$ .

- $*$  is *commutative* if  $a * b = b * a$  for all  $a, b \in A$ .
- $*$  is *associative* if  $(a * b) * c = a * (b * c)$ .

## EXAMPLE

- 1 Multiplication and addition give operators on  $\mathbb{Z}$  which are both commutative and associative.
- 2 Subtraction is an operation on  $\mathbb{Z}$  which is neither commutative nor associative.
- 3 The binary operation on  $\mathbb{Z}$  given by  $x * y = 1 + xy$  is commutative but not associative. For example  $(1 * 2) * 3 = 3 * 3 = 10$  while  $1 * (2 * 3) = 1 * (7) = 8$ .

## DEFINITION

Suppose that  $*$  is a binary operation on a nonempty set  $A$  and that  $B \subseteq A$ . If it is true that  $a * b \in B$  for all  $a, b \in B$ , then we say that  $B$  is closed under  $*$ .

## EXAMPLE

Consider multiplication on  $\mathbb{Z}$ . The set of even integers is closed under addition.

## PROOF.

Suppose that  $a, b \in \mathbb{Z}$  are even.

Then there are  $x, y \in \mathbb{Z}$  such that  $a = 2x$  and  $b = 2y$ .

Thus  $a + b = 2x + 2y = 2(x + y)$  which is even.

Since  $a$  and  $b$  were arbitrary even integers, it follows that the set of even integers is closed under addition. □

## DEFINITION

Let  $*$  be a binary operation on a nonempty set  $A$ . An element  $e$  is called an identity element with respect to  $*$  if

$$e * x = x = x * e$$

for all  $x \in A$ .

## EXAMPLE

- 1 is an identity element for multiplication on the integers.
- 0 is an identity element for addition on the integers.
- If  $*$  is defined on  $\mathbb{Z}$  by  $x * y = x + y + 1$  Then -1 is the identity.
- The operation  $*$  defined on  $\mathbb{Z}$  by  $x * y = 1 + xy$  has no identity element.

# RIGHT, LEFT AND TWO-SIDED INVERSES

## DEFINITION

Suppose that  $*$  is a binary operation on a nonempty set  $A$  and that  $e$  is an identity element with respect to  $*$ . Suppose that  $a \in A$ .

- If there exists  $b \in A$  such that  $a * b = e$  then  $b$  is called a *right inverse* of  $a$  with respect to  $*$ .
- If there exists  $b \in A$  such that  $b * a = e$  then  $b$  is called a *left inverse* of  $a$  with respect to  $*$ .
- If  $b \in A$  is both a right and left inverse of  $a$  with respect to  $*$  then we simply say that  $b$  is an *inverse* of  $a$  and we say that  $a$  is *invertible*.

## EXAMPLE

- 1 Consider the operation of addition on the integers. For any integer  $a$ , the inverse of  $a$  with respect to addition is  $-a$ .
- 2 Consider the operation of multiplication on  $\mathbb{Z}$ . The invertible elements are 1 and -1.

## FACT

*Suppose that  $*$  is a binary operation on a nonempty set  $A$ . If there is an identity element with respect to  $*$  then it is unique. In the case that there is an identity element and that  $*$  is associative then for each  $a \in A$  if there is an inverse of  $a$  then it is unique.*