

# MTHSC 412 SECTION 3.1 –GROUPS

Kevin James

## DEFINITION

A *group* is a set  $G$  together with a binary operation  $*$  on  $G$  satisfying the following conditions.

- 1  $*$  is associative
- 2  $G$  has an identity element with respect to  $*$ .
- 3 For each  $g \in G$  there is an inverse  $g^{-1}$  of  $g$  with respect to  $*$ .

## DEFINITION

A *group* is a set  $G$  together with a binary operation  $*$  on  $G$  satisfying the following conditions.

- 1  $*$  is associative
- 2  $G$  has an identity element with respect to  $*$ .
- 3 For each  $g \in G$  there is an inverse  $g^{-1}$  of  $g$  with respect to  $*$ .

## DEFINITION

Suppose that  $G$  is a group with respect to  $*$ . Then  $G$  is an *abelian* or *commutative* group if  $*$  is commutative.

## EXAMPLE

- 1 If  $A$  is any set then  $\mathcal{S}(A)$  is a group with respect to composition of functions. This group is not abelian.

## EXAMPLE

- ① If  $A$  is any set then  $\mathcal{S}(A)$  is a group with respect to composition of functions. This group is not abelian.
- ②  $\mathbb{Z}$  is an abelian group under addition.

## EXAMPLE

- ❶ If  $A$  is any set then  $\mathcal{S}(A)$  is a group with respect to composition of functions. This group is not abelian.
- ❷  $\mathbb{Z}$  is an abelian group under addition.
- ❸  $M_n(\mathbb{R})$  is an abelian group under addition.

## EXAMPLE

- ❶ If  $A$  is any set then  $\mathcal{S}(A)$  is a group with respect to composition of functions. This group is not abelian.
- ❷  $\mathbb{Z}$  is an abelian group under addition.
- ❸  $M_n(\mathbb{R})$  is an abelian group under addition.
- ❹  $\mathbb{Z}_n$  is an abelian group under addition.

## NOTATION (CYCLE NOTATION FOR $S_n$ )

- 1 Recall that if  $A = \{1, 2, \dots, n\}$  then we denote by  $S_n = \mathcal{S}(A)$ .



## NOTATION (CYCLE NOTATION FOR $S_n$ )

- 1 Recall that if  $A = \{1, 2, \dots, n\}$  then we denote by  $S_n = \mathcal{S}(A)$ .
- 2 We shall use the notation  $\sigma = (a_1, a_2, \dots, a_k)$  to denote the permutation  $\sigma \in S_n$  defined by

$$\sigma(m) = \begin{cases} a_{i+1} & \text{if } m = a_i \text{ where } 1 \leq i \leq k-1, \\ a_1 & \text{if } m = a_k. \\ m & \text{otherwise.} \end{cases}$$

## NOTATION (CYCLE NOTATION FOR $S_n$ )

- 1 Recall that if  $A = \{1, 2, \dots, n\}$  then we denote by  $S_n = \mathcal{S}(A)$ .
- 2 We shall use the notation  $\sigma = (a_1, a_2, \dots, a_k)$  to denote the permutation  $\sigma \in S_n$  defined by

$$\sigma(m) = \begin{cases} a_{i+1} & \text{if } m = a_i \text{ where } 1 \leq i \leq k-1, \\ a_1 & \text{if } m = a_k. \\ m & \text{otherwise.} \end{cases}$$

## EXAMPLE

As an element of  $S_3$ ,  $f = (1, 2)$  denotes the function  $f$  on  $\{1, 2, 3\}$  whose values are  $f(1) = 2, f(2) = 1, f(3) = 3$ .

### EXAMPLE

The elements of  $S_3$  are  $e = (1)$ ,  $(1, 2)$ ,  $(1, 3)$ ,  $(2, 3)$ ,  $(1, 2, 3)$ ,  $(1, 3, 2)$ .

### EXAMPLE

The elements of  $S_3$  are  $e = (1)$ ,  $(1, 2)$ ,  $(1, 3)$ ,  $(2, 3)$ ,  $(1, 2, 3)$ ,  $(1, 3, 2)$ .

Note that  $(1, 2, 3)^2 = (1, 3, 2)$ .

## EXAMPLE

The elements of  $S_3$  are  $e = (1)$ ,  $(1, 2)$ ,  $(1, 3)$ ,  $(2, 3)$ ,  $(1, 2, 3)$ ,  $(1, 3, 2)$ .

Note that  $(1, 2, 3)^2 = (1, 3, 2)$ .

The multiplication table for  $S_3$  is

*	(1)	(1, 2)	(1, 3)	(2, 3)	(1, 2, 3)	(1, 3, 2)
(1)	(1)	(1, 2)	(1, 3)	(2, 3)	(1, 2, 3)	(1, 3, 2)
(1, 2)	(1, 2)	(1)	(1, 3, 2)	(1, 2, 3)	(2, 3)	(1, 3)
(1, 3)	(1, 3)	(1, 2, 3)	(1)	(1, 3, 2)	(1, 2)	(2, 3)
(2, 3)	(2, 3)	(1, 3, 2)	(1, 2, 3)	(1)	(1, 3)	(1, 2)
(1, 2, 3)	(1, 2, 3)	(1, 3)	(2, 3)	(1, 2)	(1, 3, 2)	(1)
(1, 3, 2)	(1, 3, 2)	(2, 3)	(1, 2)	(1, 3)	(1)	(1, 2, 3)

### EXAMPLE

Let  $G = \{\pm 1, \pm i\}$  where  $i \in \mathbb{C}$  satisfies  $i^2 = -1$ .

### EXAMPLE

Let  $G = \{\pm 1, \pm i\}$  where  $i \in \mathbb{C}$  satisfies  $i^2 = -1$ .  
Then  $G$  is a group under multiplication of complex numbers.

## EXAMPLE

Let  $G = \{\pm 1, \pm i\}$  where  $i \in \mathbb{C}$  satisfies  $i^2 = -1$ .

Then  $G$  is a group under multiplication of complex numbers.

The multiplication table for  $G$  is

*	1	-1	$i$	$-i$
1	1	-1	$i$	$-i$
-1	-1	1	$-i$	$i$
$i$	$i$	$-i$	-1	1
$-i$	$-i$	$i$	1	-1



## DEFINITION

If a group  $G$  has a finite number of elements then we say that  $G$  has *finite order* or that  $G$  is a *finite group*.

## DEFINITION

If a group  $G$  has a finite number of elements then we say that  $G$  has *finite order* or that  $G$  is a *finite group*.

In this case, the number of elements of  $G$  is called the *order* of  $G$  and is denoted as  $o(G)$  or  $|G|$  or  $\#G$ .

## DEFINITION

If a group  $G$  has a finite number of elements then we say that  $G$  has *finite order* or that  $G$  is a *finite group*.

In this case, the number of elements of  $G$  is called the *order* of  $G$  and is denoted as  $o(G)$  or  $|G|$  or  $\#G$ .

If  $G$  does not have a finite number of elements then it is said to be an *infinite group*.