

MTHSC 412 SECTION 6.2 – QUOTIENT RINGS AND HOMOMORPHISMS

Kevin James

THEOREM

Let $I \trianglelefteq R$. If $a + I = b + I$ and $c + I = d + I$, then

$$(a + c) + I = (b + d) + I \quad \text{and} \quad ac + I = bd + I.$$

THEOREM

Let $I \trianglelefteq R$. If $a + I = b + I$ and $c + I = d + I$, then

$$(a + c) + I = (b + d) + I \quad \text{and} \quad ac + I = bd + I.$$

DEFINITION

Suppose that $I \trianglelefteq R$. Then we define addition and multiplication on R/I by

$$(a + I) + (b + I) = (a + b) + I \quad \text{and} \quad (a + I)(b + I) = ab + I.$$

EXAMPLE

Suppose that $R = \mathbb{Z}[x]$ and that $I = (x^2, 2)$. What are the distinct cosets for R/I ? Write out addition and multiplication tables for R/I .

EXAMPLE

Suppose that $R = \mathbb{Z}[x]$ and that $I = (x^2, 2)$. What are the distinct cosets for R/I ? Write out addition and multiplication tables for R/I .

THEOREM

Suppose that $I \trianglelefteq R$. Then,

- 1 R/I is a ring.
- 2 If R is commutative then R/I is commutative.
- 3 If R has an identity, then R/I has identity $1_R + I$.

DEFINITION

R/I is called the quotient ring or factor ring of R by I .

DEFINITION

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. We define the kernel of f as

$$\ker(f) = \{r \in R : f(r) = 0_S\}.$$

DEFINITION

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. We define the kernel of f as

$$\ker(f) = \{r \in R : f(r) = 0_S\}.$$

THEOREM

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. Then $\ker(f) \trianglelefteq R$.

DEFINITION

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. We define the kernel of f as

$$\ker(f) = \{r \in R : f(r) = 0_S\}.$$

THEOREM

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. Then $\ker(f) \trianglelefteq R$.

THEOREM

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. Then $\ker(f) = \{0\}$ if and only if f is injective.

THEOREM

Suppose that $I \trianglelefteq R$. Then the map $\pi : R \rightarrow R/I$ given by $\pi(r) = r + I$ is a surjective ring homomorphism with kernel I .

THEOREM

Suppose that $I \trianglelefteq R$. Then the map $\pi : R \rightarrow R/I$ given by $\pi(r) = r + I$ is a surjective ring homomorphism with kernel I .

DEFINITION

The map π is called the natural homomorphism from R to R/I .

THEOREM

Suppose that $I \trianglelefteq R$. Then the map $\pi : R \rightarrow R/I$ given by $\pi(r) = r + I$ is a surjective ring homomorphism with kernel I .

DEFINITION

The map π is called the natural homomorphism from R to R/I .

DEFINITION

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. Then we say that S is a homomorphic image of R .

THEOREM

Suppose that $I \trianglelefteq R$. Then the map $\pi : R \rightarrow R/I$ given by $\pi(r) = r + I$ is a surjective ring homomorphism with kernel I .

DEFINITION

The map π is called the natural homomorphism from R to R/I .

DEFINITION

Suppose that $f : R \rightarrow S$ is a homomorphism of rings. Then we say that S is a homomorphic image of R .

If f is an isomorphism then we say that S is an isomorphic image of R .

THEOREM (FIRST ISOMORPHISM THEOREM)

Suppose that $f : R \rightarrow S$ is a surjective homomorphism of rings with kernel K . Then $R/K \cong S$.

THEOREM (FIRST ISOMORPHISM THEOREM)

Suppose that $f : R \rightarrow S$ is a surjective homomorphism of rings with kernel K . Then $R/K \cong S$.

EXAMPLE

Consider the ring $R = \{f : \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$. Let $I = \{f \in R \mid f(2) = 0 = f(3)\}$. Describe R/I .

THEOREM (FIRST ISOMORPHISM THEOREM)

Suppose that $f : R \rightarrow S$ is a surjective homomorphism of rings with kernel K . Then $R/K \cong S$.

EXAMPLE

Consider the ring $R = \{f : \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$. Let $I = \{f \in R \mid f(2) = 0 = f(3)\}$. Describe R/I .

EXAMPLE

Describe all homomorphic images of \mathbb{Z} .