

# QUOTIENT GROUPS AND HOMOMORPHISMS: DEFINITIONS AND EXAMPLES

Kevin James

## DEFINITION

If  $\phi : G \rightarrow H$  is a homomorphism of groups, the kernel of  $\phi$  is the set

$$\ker(\phi) = \{g \in G \mid \phi(g) = 1_H\}.$$

## DEFINITION

If  $\phi : G \rightarrow H$  is a homomorphism of groups, the kernel of  $\phi$  is the set

$$\ker(\phi) = \{g \in G \mid \phi(g) = 1_H\}.$$

## PROPOSITION

Suppose that  $\phi : G \rightarrow H$  is a group homomorphism.

- 1  $\phi(1_G) = 1_H$ .
- 2  $\phi(g^{-1}) = (\phi(g))^{-1}, \forall g \in G$ .
- 3  $\phi(g^n) = \phi(g)^n, \forall g \in G$ .
- 4  $\ker(\phi) \leq G$ .
- 5  $\text{im}(\phi) \leq H$ .

## DEFINITION

Suppose that  $\phi : G \rightarrow H$  is a homomorphism of groups. For  $h \in H$ , we define the fiber above  $h$  to be  $X_h = \phi^{-1}(h)$ .

## DEFINITION

Suppose that  $\phi : G \rightarrow H$  is a homomorphism of groups. For  $h \in H$ , we define the fiber above  $h$  to be  $X_h = \phi^{-1}(h)$ .

## PROPOSITION

*Suppose that  $\phi : G \rightarrow H$  is a homomorphism of groups and let  $K = \ker(\phi)$ , we denote by  $G/K$  the set of all fibers. The set  $G/K$  is a group with operation defined by  $X_a X_b = X_{ab}$ . This group is called the quotient group of  $G$  by  $K$*

## PROPOSITION

Suppose that  $\phi : G \rightarrow H$  is a homomorphism of groups and let  $K = \ker(\phi)$ . Let  $X \in G/K$ . Then,

- 1 For any  $u \in X$ ,  $X = \{uk \mid k \in K\} := uK$ .
- 2 For any  $u \in X$ ,  $X = \{ku \mid k \in K\} := Ku$ .

## PROPOSITION

Suppose that  $\phi : G \rightarrow H$  is a homomorphism of groups and let  $K = \ker(\phi)$ . Let  $X \in G/K$ . Then,

- 1 For any  $u \in X$ ,  $X = \{uk \mid k \in K\} := uK$ .
- 2 For any  $u \in X$ ,  $X = \{ku \mid k \in K\} := Ku$ .

## DEFINITION

For any  $N \leq G$  and  $g \in G$  let

$$gN = \{gn \mid n \in N\} \quad Ng = \{ng \mid n \in N\}.$$

called respectively a left coset of  $N$  and a right coset of  $N$ .

## THEOREM

*Let  $G$  be a group and let  $K \leq G$  be the kernel of some homomorphism from  $G$  to some other group. The set of left cosets of  $K$  with operation defined by  $(uK) \cdot (wK) = uwK$  forms a group  $G/K$ . The same is true if we replace “left coset” by “right coset.”*



## THEOREM

Let  $G$  be a group and let  $K \leq G$  be the kernel of some homomorphism from  $G$  to some other group. The set of left cosets of  $K$  with operation defined by  $(uK) \cdot (wK) = uwK$  forms a group  $G/K$ . The same is true if we replace “left coset” by “right coset.”

## PROPOSITION

Let  $N \leq G$ . The set of left cosets of  $N$  in  $G$  form a partition of  $G$ . Furthermore, for all  $u, w \in G$ ,  $uN = wN$  if and only if  $w^{-1}u \in N$ .

## PROPOSITION

Let  $N \leq G$ .

- 1 The operation defined on the set of left cosets of  $N$  by  $uN \cdot wN = uwN$  is well defined if and only if  $gNg^{-1} = N$ ,  $\forall g \in G$ .
- 2 In the case that the above operation is well defined, it makes  $G/N$  the set of left cosets of  $N$  into a group.

## PROPOSITION

Let  $N \leq G$ .

- 1 The operation defined on the set of left cosets of  $N$  by  $uN \cdot wN = uwN$  is well defined if and only if  $gNg^{-1} = N$ ,  $\forall g \in G$ .
- 2 In the case that the above operation is well defined, it makes  $G/N$  the set of left cosets of  $N$  into a group.

## DEFINITION

Suppose that  $N \leq G$ . We say that  $g$  normalizes  $N$  if  $gNg^{-1} = N$ . The subgroup  $N$  is said to be normal if every element of  $G$  normalizes  $N$ . If  $N$  is a normal subgroup of  $G$ , we write  $N \trianglelefteq G$ .

## THEOREM

Let  $N \leq G$ . The following are equivalent.

- 1  $N \trianglelefteq G$ .
- 2  $N_G(N) = G$ .
- 3  $gN = Ng, \forall g \in G$ .
- 4 The operation  $(uN) \cdot (wN) = uwN$  is a well defined and makes  $G/N$  into a group.
- 5  $gNg^{-1} \subseteq N, \forall g \in G$ .

## THEOREM

Let  $N \leq G$ . The following are equivalent.

- 1  $N \trianglelefteq G$ .
- 2  $N_G(N) = G$ .
- 3  $gN = Ng, \forall g \in G$ .
- 4 The operation  $(uN) \cdot (wN) = uwN$  is a well defined and makes  $G/N$  into a group.
- 5  $gNg^{-1} \subseteq N, \forall g \in G$ .

## PROPOSITION

Suppose that  $N \leq G$ . Then,  $N \trianglelefteq G$  if and only if  $N = \ker(\phi)$  for some homomorphism  $\phi$ .

## DEFINITION

Suppose that  $N \trianglelefteq G$ . Then the homomorphism  $\pi : G \rightarrow G/N$  given by  $\pi(g) = gN$  (Check that this is indeed a homomorphism.) is called the natural projection of  $G$  onto  $G/N$ .

## DEFINITION

Suppose that  $N \trianglelefteq G$ . Then the homomorphism  $\pi : G \rightarrow G/N$  given by  $\pi(g) = gN$  (Check that this is indeed a homomorphism.) is called the natural projection of  $G$  onto  $G/N$ .

## NOTE

The natural projection of  $G$  onto  $G/N$  gives a one to one correspondence between the subgroups of  $G/N$  and the subgroups  $H$  of  $G$  satisfying  $N \leq H \leq G$ .