

# POLYNOMIAL RINGS ESPECIALLY OVER FIELDS

Kevin James

## PROPOSITION

Let  $R$  be an integral domain. Then,

- 1 If  $p(x), q(x) \in R[x]$  are nonzero, then  $\deg(p(x)q(x)) = \deg(p(x)) + \deg(q(x))$ .
- 2  $(R[x])^\times = R^\times$ .
- 3  $R[x]$  is an integral domain.

## PROPOSITION

Let  $R$  be an integral domain. Then,

- 1 If  $p(x), q(x) \in R[x]$  are nonzero, then  $\deg(p(x)q(x)) = \deg(p(x)) + \deg(q(x))$ .
- 2  $(R[x])^\times = R^\times$ .
- 3  $R[x]$  is an integral domain.

## PROPOSITION

Suppose that  $I \trianglelefteq R$  and let  $(I) = I[x]$  denote the ideal of  $R[x]$  generated by  $I$ . Then,

$$R[x]/(I) \cong (R/I)[x].$$

In particular, if  $I$  is a prime ideal of  $R$ , then  $(I)$  is a prime ideal of  $R[x]$ .

## DEFINITION

The polynomial ring in  $n$  variables with coefficients in  $R$  is defined inductively for  $n \geq 2$  as

$$R[x_1, x_2, \dots, x_n] = R[x_1, \dots, x_{n-1}][x_n].$$

## DEFINITION

The polynomial ring in  $n$  variables with coefficients in  $R$  is defined inductively for  $n \geq 2$  as

$$R[x_1, x_2, \dots, x_n] = R[x_1, \dots, x_{n-1}][x_n].$$

## NOTE

- ① For  $f \in R[x_1, x_2, \dots, x_n]$ ,  $f$  can be expressed as a finite sum of the form

$$f(x_1, x_2, \dots, x_n) = \sum_{0 \leq m_1, m_2, \dots, m_n} a_{m_1, m_2, \dots, m_n} x_1^{m_1} x_2^{m_2} \cdots x_n^{m_n},$$

where the  $a_{\vec{m}} \in R$ .

- ② We define degree as follows:

- ①  $\deg_i(a_{m_1, m_2, \dots, m_n} x_1^{m_1} x_2^{m_2} \cdots x_n^{m_n}) = m_i$ .
- ②  $\deg(a_{m_1, m_2, \dots, m_n} x_1^{m_1} x_2^{m_2} \cdots x_n^{m_n}) = \max_{1 \leq i \leq n} (\deg_i(a_{m_1, m_2, \dots, m_n} x_1^{m_1} x_2^{m_2} \cdots x_n^{m_n}))$ .
- ③  $\deg(f) = \max_{\vec{m}} (\deg(a_{\vec{m}} x^{\vec{m}}))$ .

### EXAMPLE

$$\deg(x^2y + xy^2) = 3.$$

## THEOREM

*Let  $F$  be a field. The polynomial ring  $F[x]$  is a Euclidean Domain.*

## THEOREM

*Let  $F$  be a field. The polynomial ring  $F[x]$  is a Euclidean Domain. Specifically if  $a(x), b(x) \in F[x]$  with  $0 \neq b(x)$ , then there is a unique  $q(x), r(x) \in F[x]$  such that*

$$a(x) = b(x)q(x) + r(x) \text{ with } r(x) = 0 \text{ or } \deg(r(x)) < \deg(b(x)).$$



## THEOREM

Let  $F$  be a field. The polynomial ring  $F[x]$  is a Euclidean Domain. Specifically if  $a(x), b(x) \in F[x]$  with  $0 \neq b(x)$ , then there is a unique  $q(x), r(x) \in F[x]$  such that

$$a(x) = b(x)q(x) + r(x) \text{ with } r(x) = 0 \text{ or } \deg(r(x)) < \deg(b(x)).$$

## COROLLARY

If  $F$  is a field, then  $F[x]$  is a PID and a UFD.