

**MthSc 208, Fall 2010 (Differential Equations)**

**Dr. Matthew Macauley**

**HW 3**

**Due Monday August 30th, 2010**

- (1) Suppose a cold beer at  $40^{\circ}\text{F}$  is placed into a warm room at  $70^{\circ}\text{F}$ . Suppose 10 minutes later, the temperature of the beer is  $48^{\circ}\text{F}$ . Use Newton's law of cooling to find the temperature 25 minutes after the beer was placed into the room.
- (2) A murder victim is discovered at midnight at the temperature of the body is recorded at  $31^{\circ}\text{C}$ . One hour later, the temperature of the body is  $29^{\circ}\text{C}$ . Assume that the surrounding air temperature remains constant at  $21^{\circ}\text{C}$ . Use Newton's law of cooling (the differential equation  $T' = k(A - T)$ ) to calculate the victim's time of death (when his body temperature was  $37^{\circ}\text{C}$ ).
- (3) A parachutist of mass 60 kg free-falls from an airplane at an altitude of 5000 meters. He is subjected to an air resistance force proportional to his speed. Assume that the constant of proportionality is  $r = 10$  kg/sec.
  - (a) Find and solve the differential equation governing the altitude of the parachuter at time  $t$  seconds after the start of his free-fall.
  - (b) Assuming he does not deploy his parachute, find his limiting velocity and how much time will elapse before he hits the ground (you may need to use a computer for this last part, a visual approximation from the appropriate graph is fine).
- (4) In our model of air resistance, the resistance force has depended only on the velocity. However, for an object that drops a considerable distance, such as the parachutist in the previous exercise, there is a dependence on the altitude as well. It is reasonable to assume that the resistance force is proportional to air pressure, as well as to the velocity. Furthermore, to a first-order approximation, the air pressure varies exponentially with the altitude (i.e., it is proportional to  $e^{-ax}$ , where  $a$  is a constant and  $x$  is the altitude). Propose and justify (*but do not solve!*) a differential equation model for the velocity of a falling object subject to such a resistance force.
- (5) For each of the first-order differential equations, decide whether it is linear or nonlinear. If the equation is linear, state whether it is homogeneous or inhomogeneous.
  - (a)  $y' = ky$
  - (b)  $y' = k(72 - y)$
  - (c)  $y' = y(4 - y)$
  - (d)  $y' = t + y$
  - (e)  $3y' + 5y = 3 \cos 2t$
  - (f)  $3y' + 5y = 3 \cos 2y$
  - (g)  $y' = 4t^2y - \sin t$
  - (h)  $y' = 4ty^2 - \sin t$