

Read: Strang, Section 4.3, 4.4.

Suggested short conceptual exercises: Strang, Section 4.3, #12–16, 25, 26, 29. Section 4.4, #3, 4, 8, 9, 19.

1. For this problem, consider the four data points $(t_i, b_i) = (0, 0), (1, 8), (3, 8),$ and $(4, 20)$. Let $\mathbf{t} = (0, 1, 3, 4)$ be the vector of inputs and $\mathbf{b} = (0, 8, 8, 20)$ the vector of outputs. Feel free to use a computer to solve any systems of equations you encounter throughout this problem.
 - (a) If there were a straight line $b = C + Dt$ through these four points, then a certain equation $\mathbf{A}\mathbf{x} = \mathbf{b}$ would have a solution, where $\mathbf{x} = (C, D)$. Write this equation in matrix form (that is, find \mathbf{A}).
 - (b) Instead, we wish to find the “best fit” line, which means we need to solve $\mathbf{A}\hat{\mathbf{x}} = \mathbf{p}$, where \mathbf{p} is the projection of \mathbf{b} onto the column space of \mathbf{A} . Write down the *normal equations* $\mathbf{A}^T\mathbf{A}\hat{\mathbf{x}} = \mathbf{A}^T\mathbf{b}$, where $\hat{\mathbf{x}} = (\hat{C}, \hat{D})$, and solve for $\hat{\mathbf{x}}$.
 - (c) Check that $\mathbf{e} = \mathbf{b} - \mathbf{p}$ is orthogonal to both columns of \mathbf{A} . Compute $\|\mathbf{e}\|$, which is the shortest distance from \mathbf{b} to the column space of \mathbf{A} . Sketch a diagram of $\mathbf{e}, \mathbf{b}, \mathbf{p}$, and the orthogonal subspaces $C(\mathbf{A})$ and $N(\mathbf{A}^T)$ to illustrate this.
 - (d) Plot the four data points in \mathbb{R}^2 (on the tb -plane) and sketch the best fit line through them that you just found. Clearly mark what the vectors $\mathbf{b} = (b_1, b_2, b_3, b_4), \mathbf{e} = (e_1, e_2, e_3, e_4)$, and $\mathbf{p} = (p_1, p_2, p_3, p_4)$ represent.
 - (e) Write down $E := \|\mathbf{A}\mathbf{x} - \mathbf{b}\|^2$ as a sum of four squares—the last one is $(C + 4D - 20)^2$, and compute $\partial E/\partial C$ and $\partial E/\partial D$. Set these derivatives equal to zero, and obtain scalars of the normal equations $\mathbf{A}^T\mathbf{A}\hat{\mathbf{x}} = \mathbf{A}^T\mathbf{b}$.
 - (f) The method above found the best fit degree-1 polynomial (line). Now, find the best fit degree-0 polynomial (horizontal line) $b = C$. Note that this will be a 4×1 system instead of a 4×2 system. Compute the vectors \mathbf{p} and \mathbf{e} , and the (squared) error $\|\mathbf{e}\|^2$.
 - (g) Find the best fit parabola (degree-2 polynomial) $b = C + Dt + Et^2$. On a new set of axes, plot the four data points and this parabola. Compute the vectors \mathbf{p} and \mathbf{e} , and the (squared) error $\|\mathbf{e}\|^2$.
 - (h) Find the best fit cubic (degree-3 polynomial) $b = C + Dt + Et^2 + Ft^3$. On a new set of axes, plot the four data points and this cubic. Compute the vectors \mathbf{p} and \mathbf{e} , and the (squared) error $\|\mathbf{e}\|^2$.
2. In this problem we will prove that orthonormal vectors are linearly independent two different ways.
 - (a) Vector proof: First, suppose that $c_1\mathbf{q}_1 + c_2\mathbf{q}_2 + \cdots + c_k\mathbf{q}_k = \mathbf{0}$. Show that each $c_i = 0$. [*Hint:* Start by multiplying both sides of the equation by \mathbf{q}_i^T .]
 - (b) Matrix proof: Let \mathbf{Q} be the matrix whose columns are the \mathbf{q}_i 's. Show that if $\mathbf{Q}\mathbf{x} = \mathbf{0}$, then $\mathbf{x} = \mathbf{0}$. [*Hint:* Since \mathbf{Q} need not be square, you cannot assume \mathbf{Q}^{-1} exists, but \mathbf{Q}^T of course will.]

3. For each of the following, answer either *true* (with a reason) or *false* (with a counterexample).
- If \mathbf{Q} is an orthogonal matrix, then \mathbf{Q}^{-1} is orthogonal.
 - If \mathbf{Q} is an orthogonal matrix, then \mathbf{Q}^T is orthogonal.
 - If \mathbf{Q}_1 and \mathbf{Q}_2 are orthogonal matrices, then $\mathbf{Q}_1\mathbf{Q}_2$ is orthogonal.
 - If \mathbf{Q} is a matrix with orthonormal columns (need not be square), then $\|\mathbf{Q}\mathbf{x}\| = \|\mathbf{x}\|$ for every \mathbf{x} .
4. What multiple of $\mathbf{A} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ should be subtracted from $\mathbf{b} = \begin{bmatrix} 6 \\ 0 \end{bmatrix}$ to make the resulting vector \mathbf{B} orthogonal to \mathbf{a} ? Sketch a figure showing \mathbf{A} , \mathbf{b} , and \mathbf{B} . Then normalize \mathbf{A} and \mathbf{B} to get an orthonormal set, \mathbf{q}_1 and \mathbf{q}_2 .
5. Let \mathbf{a} , \mathbf{b} , and \mathbf{c} be the (independent) column vectors of the matrix

$$\mathbf{M} = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & 5 \\ 0 & 3 & 6 \end{bmatrix}.$$

Use the Gram-Schmidt process to produce an orthonormal basis \mathbf{q}_1 , \mathbf{q}_2 , and \mathbf{q}_3 . Then write $\mathbf{M} = \mathbf{Q}\mathbf{R}$, where \mathbf{Q} is orthogonal and \mathbf{R} is upper-triangular.

6. Recall that if $\|\mathbf{u}\| = 1$, then the rank-1 matrix $\mathbf{u}\mathbf{u}^T$ is the projection matrix onto \mathbf{u} . In this case, $\mathbf{Q} = \mathbf{I} - 2\mathbf{u}\mathbf{u}^T$ is a *reflection matrix*.
- Reflecting twice across the same axis is the identity. Verify that indeed, $\mathbf{Q}^2 = \mathbf{I}$.
 - Compute $\mathbf{Q}\mathbf{u}$, and simplify this expression as much as possible.
 - Suppose \mathbf{v} is orthogonal to \mathbf{u} . Compute $\mathbf{Q}\mathbf{v}$, and simplify as much as possible.
 - Describe in plain English which subspace \mathbf{Q} is reflecting across. Your answer should involve \mathbf{u} . Include a sketch.
 - Compute the reflection matrix $\mathbf{Q}_1 = \mathbf{I} - 2\mathbf{u}_1\mathbf{u}_1^T$ where $\mathbf{u}_1 = (0, 1)$. Compute $\mathbf{Q}_1\mathbf{x}_1$, where $\mathbf{x}_1 = (a, b)$, and sketch the vectors \mathbf{u}_1 , \mathbf{x}_1 , and $\mathbf{Q}_1\mathbf{x}_1$ in the plane.
 - Compute the reflection matrix $\mathbf{Q}_2 = \mathbf{I} - 2\mathbf{u}_2\mathbf{u}_2^T$ where $\mathbf{u}_2 = (0, \sqrt{2}/2, \sqrt{2}/2)$. Compute $\mathbf{Q}_2\mathbf{x}_2$, where $\mathbf{x}_2 = (1, 1, 1)$, and sketch the vectors \mathbf{u}_2 , \mathbf{x}_2 , and $\mathbf{Q}_2\mathbf{x}_2$ in \mathbb{R}^3 .