Math 2080: Differential Equations Worksheet 7.7: The two-dimensional heat equation

NAME:

1. Consider the following initial/boundary value problem for the heat equation in a square region, where the function u(x, y, t) is defined for $0 \le x \le \pi$, $0 \le y \le \pi$ and $t \ge 0$.

$$u_t = c^2 \nabla^2 u, \qquad u(x, 0, t) = u(x, \pi, t) = u(0, y, t) = u(\pi, y, t) = 0$$
$$u(x, y, 0) = 2 \sin x \sin y + 5 \sin 2x \sin y.$$

(a) Briefly describe, and sketch, a physical situation which this models. Be sure to explain the effect of the boundary conditions and the initial condition.

(b) Assume that there is a solution of the form u(x, y, t) = f(x, y)g(t). Find u_{xx} , u_{yy} , and u_t .

(c) Plug u = fg back into the PDE and divide both sides by fg (i.e., "separate variables") to get the *eigenvalue problem*. Briefly justify why this quantity must be a constant. Call this constant λ . Write down an ODE for g(t), and a PDE for f(x, y) (the *Helmholz equation*). Include four boundary conditions for f(x, y). (d) Solve the Helmholz equation and determine λ . You may assume that f(x, y) = X(x)Y(y), then separate variables.

(e) Solve the ODE for g(t).

(f) Find the general solution of the boundary value problem. It will be a superposition (infinite sum) of solutions $u_{nm}(x, y, t) = f_{nm}(x, y)g_{nm}(t)$.

(g) Find the particular solution to the initial value problem that additionally satisfies the initial condition $u(x, y, 0) = 2 \sin x \sin y + 5 \sin 2x \sin y$.

(h) What is the steady-state solution? Give a mathematical and intuitive (physical) justification.

2. Consider the following inhomogeneous BVP for the heat equation in a square region.

$$u_t = c^2 \nabla^2 u,$$
 $u(x,0) = u(0,y) = 0,$ $u(x,\pi) = \sin x,$ $u(\pi,y) = \sin 2y.$

Without knowing the initial conditions, determine the steady-state solution. (*Hint*: If you use your result from the previous worksheet, then almost no actual work is needed on this problem.)