

Identifying CpG islands using hidden Markov models

Matthew Macauley

Department of Mathematical Sciences
Clemson University
<http://www.math.clemson.edu/~macaule/>

Math 4500, Fall 2016

CpG islands

On a DNA strand, a cytosine followed by guanine is a dinucleotide called **CpG**. The 'p' is for the *phosphate bond* between them.



Figure: CpG nucleotides on a DNA strand and its complement.

CpG's are often clustered in regions called **CpG islands** (CGIs).

CGIs are often associated with the promoter region of genes (where transcription begins).

Identifying CGIs can help identify new genes, some of which may be involved in cancer.


Goal

Given a genome of millions of base pairs, how can one identify the CpG islands?

Cytosine methylation

Almost all cells in an organism have the same DNA sequence. The difference lies in the levels of *gene expression*.

One common way that genes are turned off is by a chemical change called **methylation** at the promoter CGI.



Promoter regions of housekeeping genes are usually unmethylated.


Appropriate methylation of CGIs is needed for normal development. If methylation occurs when it should not in tumor suppressor genes, then problems such as cancer can result.

In mammals, 70–80% of CpG cytosines are methylated, but it depends on the type of cell.

For example, hemoglobin genes should be methylated (and shut off) in skin cells but unmethylated (and expressed) in red blood cell precursors.

Methylation and deamination

5-methyl cytosine can be **deaminated** to produce thymine (T), which is a mutation. As a result, there is a lack of CpG sites in methylated DNA.



Rule of thumb

On an evolutionary timescale, **unmethylated C's tend to persist** and **methylated C's tend to be eliminated**.

CATTCCGCCTTCTCTCCCGAGGTGGCGCGTGGGA
 GGTGTTTTCCTCGGGTTCGTAAAGAATAGGCCAGG
 CAGCTTCCCGCGGGATGCCCTCATCCCTCTCGG
 GGTTCCGCTCCCACCGCCCGCGGTTCCGCCGTT
 CCGCCTGCGAGATGTTTTCCAGCGACAATGATTC
 CACTCTCGCGCCTCCCATGTTGATCCAGCTCCT
 CTGCGGGGTCAGGACCCCTGGGCCCGCCCG
 CTCCACTCAGTCAATCTTTGTCCCGGTATAAGGCG
 GATTATCGGGGTGGTGGGGCGGCTGATTCGSA
 CGAATGCCCTTGGGGGTACCCCGGGAGGGAACTC
 CGGGCTCGCGGCTTTGGCCACCGCCACCCCTGGT
 TGAGCGCGCCCGAGGGCACACAGGGGGCGCTCG
 ATGTTCTGACGCCCCCGCAGCAGCCCCACTCC
 CCGGCTCACCCCTCGATTGGCTGGCCCGCCCGAG
 CTCTGTGCTGTGATTGGTCACAGCCCGTGTCCGTC
 CGCGGGCGCGGGGGCGGATCGAGGTGACCGCGCA
 GAGGCCAGCTCGGGCGGTGTCCCGCCCGCGC
 GACTCGGGCGGAGTTTCGCGAGGGCGAAGCG
 GGGCAGTGTGACCGCAGCGGTCTGGGAGGCGC
 CCGCGCGCGTCCGAGCAGCTCCCGTCTCTCCCA
 GCCGTACCCCGCGCGTCCCGCGCCCTGGCC
 TCCCGCACTCGCGCACTCCTGTCCCGCCACC
 CGCCACCTCCCACCTCGATCGGTGCGCGGGCTGC
 TGCGTGATGGGCTGCGGAGCGCGCCCTGCGG
 CTCGCGCGGGGCGCGTGTCTCGCGCTGAGGTGCGT
 CGGTGCCCGGCCCCCGCGCCCGCGCGCGCGC
 GGCTCCTGTTGACCGGTCCCGCGTCCGCTGTGC
 AGCGCGCGCTGAGGTAAGCGCGCGGGCTGGCG
 CGGTTGGCGCGCGCGGTCCCGGGGTTGGGGAGGG
 GGCGCTTCCCGCGGGAGGAGCGCGCGGCCGG
 GGTCGCGCGGGGTCTGAGGGGA

CTCTTAGTTTTGGGTGCATTTGTCTGGCTTCCAAA
 CTAGATTGAAAGCTCTGAAAAAAAACCTATCTTGT
 GTTCTACTCTGTTAGCTCATAGTAGGTACCAGGA
 AGTAGTAGGGTTGACTGCATTGATTTGGGACTACAC
 TGGGAGTTTTCTTCCCATCTCCCTTAGTTTTCTCT
 TTTTTCTTTCTTTCTTTCTTTTTTTTTTTTTTTTT
 TTGAGATGTCTCTTGTCTCAGTCCCCCAGGCTGGA
 GTGCAGTGGTGCGATCTTGGCTCACTGTAGCCCTCC
 ACCTCCAGGTTCAAGCAATTCTACTGCTTAGCCT
 CCGAGTAGTGTGGGATTACAAGCACC CGCCACCAT
 TCCTGGCTAATTTTTTTTTTTTGTATTTTAGTTGAGA
 CAGGGTTTTACCATGTTGGTGTGCTGTCTCAGA
 CTCCTGGGGCCTAGCGATCCCCCTGCCTCAGCCT
 CCCAGAGTGTTAGGATTACAGGCATGAGCCACTGT
 ACCCGCCCTCTCTCCAGTTCACAGTTTGAATCCAA
 GGGAASTAAGTTTAAGATAAAGTTACGATTTTGAAT
 CTTTGGATTGAGAAGAAATTTGTACCTTTAACACCT
 AGAGTTGAACGTTTCATACCTGGAGGCCTTAACATT
 AAGCCCTAGCCAGCCTCCAGCAAGTGGACATTGGT
 CAGGTTTGGCAGGATTCTCCCTGGAAGTGGACT
 GAGAGCCACACCCTGGCCTGTCCACCATACCCATCC
 CCTATCCTTAGTGAAGCAAACTCCTTTGTTCCCTT
 CTCCTTCTCCTAGTGACAGGAAATTTGTATCCTA
 AAGAATGAAAATAGTCTACGCAACTCGTGGCTCAG
 GCCTCTTGACTTCAGGCGTCTGTTTAATCAAGT
 GACATCTTCCCGAGGCTCCCTGAATGTGGGAGATG
 AAAGAGACTAGTTCAACCCTGACTGAGGGGAAAG
 CCTTTGTGAAGGGTCAGGAG

Left: CpG sites at 1/10 nucleotides, constituting a CpG island. The sample is of a gene-promoter, the highlighted ATG constitutes the start codon.

Right: CpG sites present at every 1/100 nucleotides, constituting a more normal example of the genome, or a region of the genome that is commonly methylated.

How to define a CpG island

The human genome has a 42% GC content. Thus, the expected frequency of a CpG $0.21 \cdot 0.21 = 4.41\%$. However, the actual frequency is 1%.

The **percent combined C + G content** ($\%C + G$) is defined “exactly how you would expect.”

If dinucleotides were formed by randomly choosing two nucleotides, then the expected number of CpG's would be

$$\frac{(\# \text{ C's}) \cdot (\# \text{ G's})}{\text{length of sequence}}$$

The **observed over expected CpG ratio** (O/E CpG) is: $\frac{\text{observed } \# \text{ CpG's}}{\text{expected } \# \text{ CpG's}}$

Definition (Gardiner-Garden, Frommer, 1987)


A subsequence in a vertebrate genome is **CpG island** if:

1. it has length at least 200 bp;
2. $\%C + G \geq 50\%$;
3. $\text{O/E CpG} \geq 0.6$;

There is no universal standard for these values. Another paper (Takai & Jones) used 500 bp, $\%C + G \geq 55\%$, and $\text{O/E CpG} \geq 0.65$.

Finding CpG islands

One method for inferring CpG islands is purely algorithmic: using a **sliding window**.




The remainder of this lecture will focus on an alternative approach: **hidden Markov models**.

The occasionally dishonest casino

Suppose a casino hosts a simple game with two dice: one fair and one unfair.

- FAIR: $p(1) = p(2) = p(3) = p(4) = p(5) = p(6) = 1/6$.
- UNFAIR: $p(1) = p(2) = p(3) = p(4) = p(5) = 1/10, p(6) = 1/2$.

The casino switches between fair and unfair die according to the following probabilities:



You cannot tell which die the casino is using. This is a **hidden Markov model** (HMM).

Suppose that the outcome of the game is the following:

- WIN: roll 1, 2, 3, or 4.
- LOSE: roll 5 or 6.


Would you play this game?

Two examples of Hidden Markov models

The parameters of an HMM can be encoded in a table.


HMM for the occasionally dishonest casino

State	Transitions		Emissions		Initial distribution
	F	U	W	L	
F	.95	.05	2/3	1/3	.5
U	.1	.9	.4	.6	.5




HMM for CpG islands (simple)

State	Transitions		Emissions				Init. dist.
	-	+	A	C	T	G	
-	.95	.05	.27	.24	.26	.23	.5
+	.1	.9	.15	.33	.16	.36	.5



A better hidden Markov model for CpG islands

A “better” HMM model should incorporate the fact that transmission probabilities within CpG islands are much different than the rest of the genome.



The following is from a sequence of annotated human DNA of length $\approx 60,000$.

	Transitions								Emissions				Init.
	A ₋	C ₋	T ₋	G ₋	A ₊	C ₊	T ₊	G ₊	A	C	T	G	
A ₋	.300	.205	.210	.285	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	1	0	0	0	.125
C ₋	.322	.298	.302	.078	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	0	1	0	0	.125
T ₋	.248	.246	.208	.298	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	0	0	1	0	.125
G ₋	.177	.239	.292	.292	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	$(1-q)/4$	0	0	0	1	.125
A ₊	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$.180	.274	.120	.426	1	0	0	0	.125
C ₊	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$.171	.368	.188	.274	0	1	0	0	.125
T ₊	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$.161	.339	.125	.375	0	0	1	0	.125
G ₊	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$	$(1-p)/4$.079	.355	.182	.384	0	0	0	1	.125

Three canonical HMM problems, formalized

Problem #1: Evaluation

Given an observed path $x = x_1x_2x_3 \cdots x_\ell$, what is its probability $P(x)$? That is, compute

$$P(x) = \sum_{\pi} P(x, \pi), \quad \text{where } P(x, \pi) = a_{0\pi_1} \prod_{i=1}^{\ell} e_{\pi_i}(x_i) a_{\pi_i, \pi_{i+1}}$$

and the sum is over all hidden sequences $\pi = \pi_1\pi_2 \cdots \pi_\ell$.

Problem #2: Decoding

Given an observed path $x = x_1x_2x_3 \cdots x_\ell$, what is the most likely hidden path $\pi = \pi_1\pi_2\pi_3 \cdots \pi_\ell$ to emit x ? That is, compute

$$\pi_{\max} = \arg \max_{\pi} P(\pi|x) = \arg \max_{\pi} P(x, \pi)$$

Problem #3: Learning

Given an observed sequence x (or set of sequences), what are the HMM parameters that make x mostly likely to occur?