- 1. Let K be a finite field. The *characteristic* of K, denoted char K, is the smallest positive integer n for which  $n1 := \underbrace{1+1+\cdots+1}_{K} = 0$ .
  - (a) Prove that the characteristic of K is prime.
  - (b) Show that K is a vector space over  $\mathbb{Z}_p$ , where  $p = \operatorname{char} K$ .
  - (c) Show that the order |K| of K (the number of elements it contains) is a prime power.
  - (d) Show that if K and L are finite fields with  $K \subset L$  and  $|K| = p^m$  and  $|L| = p^n$ , then m divides n.
- 2. Let X be a vector space over a field K and let X' be the set of linear functions from X to K, also known as the *dual space* of X.
  - (a) Let  $v_1, \ldots, v_n$  be a basis for X. For each *i*, show there exists a unique linear map  $f_i: X \to K$  such that  $f_i(v_i) = 1$  and  $f_i(v_j) = 0$  for  $j \neq i$ .
  - (b) Show that  $f_1, \ldots, f_n$  is a basis for X' (called the *dual basis* of  $v_1, \ldots, v_n$ ).
  - (c) Consider the basis  $v_1 = (1, -1, 3)$ ,  $v_2 = (0, 1, -1)$ , and  $v_3 = (0, 3, -2)$  of  $X = \mathbb{R}^3$ . Find a formula for each element of the dual basis.
  - (d) Express the linear map  $f \in X'$ , where f(x, y, z) = 2x y + 3z as a linear combination of the dual basis,  $f_1, f_2, f_3$ .
- 3. Let S be a subset of X. The annihilator of S is the set

$$S^{\perp} = \{ \ell \in X' \mid \ell(s) = 0 \text{ for all } s \in S \}.$$

- (a) Show that  $S^{\perp}$  is a subspace of X'.
- (b) Show that span(S) is the intersection of all subspaces  $T_{\alpha}$  of X that contain S:

$$\operatorname{span}(S) = \bigcap_{S \subseteq T_{\alpha} \le X} T_{\alpha}$$

making it well-founded to speak of the "smallest subpace of X that contains S."

- (c) Let  $Y = \operatorname{span}(S)$ . Show that  $S^{\perp} = Y^{\perp}$ .
- 4. Let  $\mathcal{P}_2$  be the vector space of all polynomials  $p(x) = a_0 + a_1 x + a_2 x^2$  over  $\mathbb{R}$ , with degree  $\leq 2$ . Let  $\xi_1, \xi_2, \xi_3$  be distinct real numbers, and define

$$\ell_j \colon \mathcal{P}_2 \longrightarrow \mathbb{R}, \qquad \ell_j(p) = p(\xi_j) \quad \text{for} \quad j = 1, 2, 3.$$

- (a) Show that  $\ell_1, \ell_2, \ell_3$  is a basis for the dual space  $\mathcal{P}'_2$ .
- (b) Find polynomials  $p_1(x), p_2(x), p_3(x)$  in  $\mathcal{P}_2$  of which  $\ell_1, \ell_2, \ell_3$  is the dual basis in  $\mathcal{P}'_2$ .
- 5. Let W be the subspace of  $\mathbb{R}^4$  spanned by (1, 0, -1, 2) and (2, 3, 1, 1). Which linear functions  $\ell(x) = c_1 x_1 + c_2 x_2 + c_3 x_3 + c_4 x_4$  are in the annihilator of W? Write your answer by giving an explicit basis of  $W^{\perp}$ .