

## Math 2080: Differential Equations

### Worksheet 4.7: Phase portraits with repeated eigenvalues

NAME:

Consider the system of differential equations:  $\begin{cases} x_1' = 4x_1 + x_2, & x_1(0) = -1 \\ x_2' = -1x_1 + 2x_2, & x_2(0) = 1 \end{cases}$

- (a) Write this in matrix form,  $\mathbf{x}' = \mathbf{A}\mathbf{x}$ ,  $\mathbf{x}(0) = \mathbf{x}_0$ .
- (b) Knowing that  $\mathbf{A}$  has a repeated eigenvalue,  $\lambda_{1,2} = 3$ , and only one eigenvector,  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ , write down a solution  $\mathbf{x}_1(t)$  to  $\mathbf{x}' = \mathbf{A}\mathbf{x}$ .
- (c) To find a second solution, assume that  $\mathbf{x}_2(t) = te^{3t}\mathbf{v} + e^{3t}\mathbf{w}$ . Plug this back into  $\mathbf{x}' = \mathbf{A}\mathbf{x}$  and equate coefficients (of  $te^{3t}$  and  $e^{3t}$ ) to get a system of two equations, involving  $\mathbf{v}$ ,  $\mathbf{w}$ , and  $\mathbf{A}$ .

- (d) Solve for  $\mathbf{v}$  by inspection. Plug this back into the second equation and solve for  $\mathbf{w}$  (it will involve a constant,  $C$ ).
- (e) Using what you got for  $\mathbf{v}(t)$  and  $\mathbf{w}(t)$ , write down a solution  $\mathbf{x}_2(t)$  that is not a scalar multiple of  $\mathbf{x}_1$ . (Pick the simplest value of  $C$  that works.)
- (f) Write down the general solution,  $\mathbf{x}(t)$ . As  $t \rightarrow \infty$ , which of the three terms of  $\mathbf{x}(t)$  “grows faster”?
- (g) Sketch the phase portrait. To determine which way the curves “spiral”, compute  $\mathbf{x}' = \mathbf{A}\mathbf{x}$  at  $\mathbf{x} = \begin{bmatrix} 10 \\ 0 \end{bmatrix}$  and see if this velocity vector is pointing upwards or downwards.