


Read the following, which can all be found either in the textbook or on the course website.

- Chapters 6 of *Visual Group Theory*, or Chapters 4.1, 5.4, 5.5, 7 of *IBL Abstract Algebra*.
- VGT Exercises 6.6–6.9, 6.12, 6.17–6.20, 6.28–6.30.

Write up solutions to the following exercises.

1. A Cayley diagram and multiplication table for the dihedral group D_4 are shown below.



Section 2 of the class lecture notes describes two algorithms for expressing a group G of order n as a set of permutations in S_n . One algorithm uses the Cayley diagram and the other uses the multiplication table. In this problem, you will explore this a bit further.

- (a) Label the vertices of the Cayley diagram from the set $\{1, \dots, 8\}$ and use this to construct a permutation group isomorphic to D_4 , and sitting inside S_8 .
 - (b) Label the entries of the multiplication table from the set $\{1, \dots, 8\}$ and use this to construct a permutation group isomorphic to D_4 , and sitting inside S_8 .
 - (c) Are the two groups you got in Parts (a) and (b) the same? (The answer will depend on your choice of labeling.) If “yes”, then repeat Part (a) with a different labeling to yield a different group. If “no”, then repeat Part (a) with a different labeling to yield the group you got in Part (b).
2. Find all subgroups of the following groups, and arrange them in a Hasse diagram, or subgroup lattice. Moreover, label each edge between $K \leq H$ with the index, $[H : K]$.
 - (a) $C_{23} = \langle r \mid r^{23} = 1 \rangle$;
 - (b) $C_{24} = \langle r \mid r^{24} = 1 \rangle$;
 - (c) $\mathbb{Z}_3 \times \mathbb{Z}_3 = \{(a, b) \mid a, b \in \{0, 1, 2\}\}$;
 - (d) $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 = \{(a, b, c) \mid a, b, c \in \{0, 1\}\}$; (*Tip*: it’s notationally easier to write elements as binary strings, e.g., abc instead of (a, b, c));
 - (e) $S_3 = \{e, (1\ 2), (2\ 3), (1\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$;
 - (f) $Q_8 = \langle i, j, k \mid i^2 = j^2 = k^2 = ijk = -1 \rangle$.

3. For each subgroup H of S_4 described below, write out all of its elements and determine what well-known group it is isomorphic to.

- (a) $H = \langle (1\ 2), (3\ 4) \rangle$;
- (b) $H = \langle (1\ 2)(3\ 4), (1\ 3)(2\ 4) \rangle$;
- (c) $H = \langle (1\ 2), (2\ 3) \rangle$;
- (d) $H = \langle (1\ 2), (1\ 3\ 2\ 4) \rangle$;
- (e) $H = \langle (1\ 2\ 3), (2\ 3\ 4) \rangle$.

4. Prove the following, algebraically (that is, do not refer to Cayley diagrams):

- (a) If \mathcal{H} is a collection of subgroups of G , then the intersection $\bigcap_{H \in \mathcal{H}} H$ is also a subgroup of G .
- (b) For any (possibly infinite) subset $S \subseteq G$, the subgroup generated by S is defined as

$$\langle S \rangle := \{s_1^{e_1} s_2^{e_2} \cdots s_k^{e_k} \mid s_i \in S, e_i \in \{-1, 1\}\}.$$

That is, $\langle S \rangle$ consists of all finite “words” that can be written using the elements in S and their inverses. Note that the s_i ’s need not be distinct. Prove that

$$\langle S \rangle = \bigcap_{S \subseteq H \leq G} H,$$

where the intersection is taken over all subgroups of G that contain S . [*Hint*: One way to prove that $A = B$ is to show that $A \subseteq B$ and $B \subseteq A$.]

5. For a subgroup $H \leq G$ and element $x \in G$, the set $xH := \{xh \mid h \in H\}$ is a *left coset* of H .

- (a) Prove that if $x \in H$, then $xH = H$. What is the interpretation of this statement in terms of the Cayley diagram?
- (b) Prove that if $b \in aH$, then $aH = bH$.
- (c) Prove that all left cosets have the same size. One way to do this is to prove that for any $x \in G$, the map

$$\varphi: H \longrightarrow xH, \quad \varphi: h \longmapsto xh$$

is a bijection.

- (d) Conclude that G is partitioned by the left cosets of H , all of which are equal size.

6. A subgroup H of G is *normal* if $xH = Hx$ for all $x \in G$. Prove that if $[G : H] = 2$, then H is a normal subgroup of G . [*Hint*: Use the results of the previous problem.]