

Read Chapters 10.6–10.7 of *Visual Group Theory* or Chapter 22 of *AATA*. Then write up solutions to the following exercises.

1. The splitting field of $f(x) = x^4 - 3$ is $K = \mathbb{Q}(\sqrt[4]{3}, i)$. Since this is a degree-8 extension over \mathbb{Q} (see HW 11), its Galois group has order 8.
 - (i) In the complex plane, sketch the roots of $f(x)$, and all 4th roots of unity: $\pm 1, \pm i$.
 - (ii) Compute the Galois group of $f(x)$. Write down two automorphisms, r and f , that generate it. It suffices to specify where they send the generators $\sqrt[4]{3}$ and i .
 - (iii) Draw the subgroup lattice of G . Each subgroup should be expressed by its generators, rather than what subgroup it is isomorphic to. Label the edges by index, and circle the subgroups that are normal in G .
 - (iv) Draw the subfield lattice of K . Label the edges by degree, and circle the subfields that are normal extensions of \mathbb{Q} . [*Hint*: The two subfields that are “easiest” to overlook are $\mathbb{Q}((1+i)\sqrt[4]{3})$ and $\mathbb{Q}((1-i)\sqrt[4]{3})$.]
 - (v) For each intermediate subfield $\mathbb{Q} \subseteq F \subseteq K$, write down the largest subgroup of G that fixes F .
 - (vi) For each subgroup $H \leq G$, write down the largest intermediate subfield fixed by H .
 - (vii) For each normal extension F of \mathbb{Q} , find a polynomial $g(x)$ whose splitting field is F .
 - (viii) For each non-normal extension E of \mathbb{Q} , find a polynomial that has one, but not all, of its roots in E .

2. The roots of $f(x) = x^n - 1$ are the n complex numbers $C_n := \{e^{2k\pi i/n} : k = 0, 1, \dots, n-1\}$, and are called the n^{th} roots of unity. A primitive root of unity is $\zeta = e^{2k\pi i/n}$ for which $\gcd(n, k) = 1$. It is easy to see that $\mathbb{Q}(\zeta)$ is the splitting field of $x^n - 1$.
 - (a) For each $n = 3, \dots, 8$, sketch the n^{th} roots of unity in the complex plane. Use a different set of axes for each n . Next to each root, write its order, as an element of C_n . Make it clear (e.g., star, or draw darker) which are the primitive roots of unity.
 - (b) Prove that for any primitive root of unity $\zeta_k = e^{2k\pi i/n}$, the mapping

$$\phi_k : \mathbb{Q}(\zeta) \rightarrow \mathbb{Q}(\zeta), \quad \phi_k(\zeta) = \zeta^k$$

is a field automorphism. That is, prove that ϕ_k is a surjective field homomorphism (every nonzero field homomorphism is injective). Why is this not an automorphism if $\gcd(n, k) \neq 1$?

- (c) Make a multiplication table of $\text{Gal}(x^n - 1)$ for $n = 3, \dots, 8$.
- (d) Describe the group $\text{Gal}(x^n - 1)$. This is a class of groups that we have previously encountered.

3. For each of the following polynomials, determine if it is irreducible. If it is not, then factor it into irreducible factors.

- (a) $f(x) = x^4 - 10x^3 + 12x^2 - 8x + 6$ over \mathbb{Q} .
- (b) $f(x) = x^4 + x^3 + x^2 + x + 1$ [Hint: Let $u = x + 1$, and change variables.]
- (c) $f(x) = x^5 - 1$ over \mathbb{Q} .
- (d) $f(x) = x^6 - 1$ over \mathbb{Q} . [Hint: Google “cyclotomic polynomial.”]
- (e) $f(x) = x^8 - 1$ over \mathbb{Q} .
- (f) $f(x) = x^{12} - 1$ over \mathbb{Q} .
- (g) $f(x) = x^3 + x^2 + x + 1$ over \mathbb{Z}_2 .
- (h) $f(x) = x^3 + x + 2$ over \mathbb{Z}_3 .


4. The splitting field of $f(x) = x^8 - 2$ over \mathbb{Q} is $\mathbb{Q}(\sqrt[8]{2}, \zeta) = \mathbb{Q}(\sqrt[8]{2}, i)$, where $\zeta = e^{2\pi i/8} = \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i$, a primitive 8th root of unity. The Galois group is generated by the automorphisms

$$\left\{ \begin{array}{l} \sigma: \sqrt[8]{2} \mapsto \zeta \sqrt[8]{2} \\ \sigma: \zeta \mapsto \zeta \end{array} \right. \quad \left\{ \begin{array}{l} \tau: \sqrt[8]{2} \mapsto \sqrt[8]{2} \\ \tau: i \mapsto -i. \end{array} \right.$$

It is isomorphic to the *quasidihedral group*, with group presentation

$$QD_8 = \langle \sigma, \tau \mid \sigma^8 = 1, \tau^2 = 1, \tau\sigma\tau = \sigma^3 \rangle,$$

and subgroup lattice shown below.



- (a) Draw Cayley diagrams of both QD_8 and the regular dihedral group, D_8 .
- (b) Sketch the roots of $f(x) = x^8 - 2$ on the complex plane, along with all 8th roots of unity: $\zeta^0, \zeta^1, \dots, \zeta^7$.
- (c) For each subgroup $H \leq \text{Gal}(x^8 - 2)$, find the largest subgroup of $\mathbb{Q}(\sqrt[8]{2}, i)$ fixed by H , and write it in the corresponding place on the subfield lattice on the right.
It is helpful to know that the proper subfields of $\mathbb{Q}(\sqrt[8]{2}, i)$ are: \mathbb{Q} , $\mathbb{Q}(i)$, $\mathbb{Q}(\sqrt{2})$, $\mathbb{Q}(\sqrt[4]{2})$, $\mathbb{Q}(\sqrt[8]{2})$, $\mathbb{Q}(\sqrt{2}i)$, $\mathbb{Q}(\sqrt[4]{2}i)$, $\mathbb{Q}(\sqrt[8]{2}i)$, $\mathbb{Q}(\sqrt{2}, i)$, $\mathbb{Q}(\sqrt[4]{2}, i)$, $\mathbb{Q}((1+i)\sqrt[4]{2})$, $\mathbb{Q}((1-i)\sqrt[4]{2})$, $\mathbb{Q}(\zeta\sqrt[8]{2})$, $\mathbb{Q}(\zeta^3\sqrt[8]{2})$.
- (d) Circle each subfield E that is a normal extension of \mathbb{Q} , and find a polynomial whose splitting field over \mathbb{Q} is E .