

Lecture 7.4: Polar decomposition

Matthew Macauley

School of Mathematical & Statistical Sciences
Clemson University
<http://www.math.clemson.edu/~macaule/>

Math 8530, Advanced Linear Algebra

The idea of the polar decomposition

Every nonzero complex number $z \in \mathbb{C}$ has a unique **polar form**

$$z = re^{i\theta} = |z|e^{i\theta}, \quad r \in \mathbb{R}^+, \quad \theta \in [0, 2\pi).$$

This can be thought of as decomposing $z \in \mathbb{C}$ into:

- a rotation by θ ,
- a scaling by $|z| = r = \sqrt{\bar{z}z}$.

This is simply the **polar decomposition** of a 1×1 matrix.

Every linear map $A \in \text{Hom}(X, X)$ can be decomposed as $A = UP$, where

- U is unitary; i.e., an **isometry** of X ,
- $P \geq 0$; a **scaling** along an orthonormal axis u_1, \dots, u_n .

It turns out that $P = \sqrt{A^*A} := |A|$, and so sometimes this is written $A = U|A|$.

In this lecture, we will derive the polar decomposition of a linear map

$$A: X \longrightarrow U, \quad \dim X = m, \quad \dim U = n.$$

In the next lecture, we will derive the celebrated **singular value decomposition (SVD)**.

Singular values

Key properties (Proposition 7.2)

- $A^*A \geq 0$;
- Every $P \geq 0$ has a **unique nonnegative square root** $R := \sqrt{P}$, such that $R^2 = P$.

This means that for some $\lambda_1, \dots, \lambda_m \geq 0$,

$$A^*A = W \begin{bmatrix} \lambda_1^2 & & \\ & \ddots & \\ & & \lambda_m^2 \end{bmatrix} W^*, \quad \text{and} \quad \sqrt{A^*A} = W \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_m \end{bmatrix} W^*.$$

Definition

The eigenvalues of $\lambda_1, \dots, \lambda_m$ of $\sqrt{A^*A}$ are called the **singular values** of A .

Facts (that we've seen)

- $\|Ax\| = \|\sqrt{A^*A}x\|$ for all $x \in X$.
- A , A^*A , and $\sqrt{A^*A}$ have the same nullspace.
- A , A^*A , and $\sqrt{A^*A}$ have the same rank.

Polar decomposition of an invertible map

Theorem

Every linear map $A: X \rightarrow U$ can be written as $A = UP$ where $P \geq 0$ and U is unitary. This is called the (left) **polar decomposition** of A .

To construct the polar decomposition, suppose $A = UP$.

Since $P \geq 0$, we can write $P = QDQ^*$, and so

$$P^*P = (QDQ^*)^*(QDQ^*) = (QD^*Q^*)QDQ^* = QD^2Q^* = P^2.$$

Now, notice that

$$A^*A = (UP)^*(UP) = P^*U^*UP = P^*P = P^2.$$

Therefore, $P = \sqrt{A^*A}$.

If A is invertible, then $U = AP^{-1} = A\sqrt{A^*A}^{-1}$ is uniquely determined.

In this case,

$$A = UP = (A\sqrt{A^*A}^{-1})\sqrt{A^*A}.$$

If A is not invertible, then U still exists, but is not unique.

Polar decomposition of an general linear map

Theorem

Every linear map $A: X \rightarrow U$ can be written as $A = UP$ where $P \geq 0$ and U is unitary. This is called the **polar decomposition** of A .

Suppose the eigenvalues of $\sqrt{A^*A}$ are

$$\lambda_1 \geq \dots \geq \lambda_r > \lambda_{r+1} = \dots = \lambda_m = 0,$$

and pick a set x_1, \dots, x_m of **orthonormal eigenvectors**. Then

$$\frac{1}{\lambda_1}Ax_1, \dots, \frac{1}{\lambda_r}Ax_r, x_{r+1}, \dots, x_m$$

is orthonormal. The polar decomposition is $A = UP$ where $P = \sqrt{A^*A}$ and

$$U = \left[\begin{array}{c|c|c|c|c|c} \frac{1}{\lambda_1}Ax_1 & \dots & \frac{1}{\lambda_r}Ax_r & x_{r+1} & \dots & x_m \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \end{array} \right] \left[\begin{array}{c} - & x_1^H & - \\ & \vdots & \\ - & x_m^H & - \end{array} \right].$$

Remark

If $A: X \rightarrow X$ and $r := \det P = |\det A|$, then

$$\det A = \det U \det P = e^{i\theta} \cdot r.$$