

Chapter 3: Group structure

Matthew Macauley



Department of Mathematical Sciences
Clemson University

<http://www.math.clemson.edu/~macaule/>

Math 8510, Abstract Algebra

Subgroup lattices


Let's compare the two groups of order 4:



- Proper subgroups of V_4 : $\langle h \rangle = \{e, h\}$, $\langle v \rangle = \{e, v\}$, $\langle r \rangle = \{e, r\}$, $\langle e \rangle = \{e\}$.
- Proper subgroups of C_4 : $\langle r \rangle = \{1, r, r^2, r^3\} = \langle r^3 \rangle$, $\langle r^2 \rangle = \{1, r^2\}$, $\langle 1 \rangle = \{1\}$.

It is illustrative to arrange them in a [subgroup lattice](#).

Order: 4



2




1

$C_4 = \langle r \rangle$

$\langle r^2 \rangle$


$\langle 1 \rangle$

The two groups of order 6



Here are their subgroup lattices:

Order: 6




3

2

1

$D_3 = \langle r, f \rangle$



Order: 6

3

2

1

Intersections of subgroups

Proposition (exercise)

For any collection $\{H_\alpha \mid \alpha \in A\}$ of subgroups of G , the intersection $\bigcap_{\alpha \in A} H_\alpha$ is a subgroup.

Every subset $S \subseteq G$, not necessarily finite, generates a subgroup, denoted

$$\langle S \rangle = \{s_1^{e_1} s_2^{e_2} \cdots s_k^{e_k} \mid s_i \in S, e_i = \{1, -1\}\}.$$

That is, $\langle S \rangle$ consists of **finite words** built from elements in S and their inverses.

Proposition (proof on board)

For any $S \subseteq G$, the subgroup $\langle S \rangle$ is the intersection of all subgroups containing S :







$$\langle S \rangle = \bigcap_{S \subseteq H_\alpha \leq G} H_\alpha,$$

That is, the subgroup **generated by S** is the **smallest subgroup containing S** .


- LHS: the subgroup built “**from the bottom up**”
- RHS: the subgroup built “**from the top down**”

There are a number of mathematical objects that can be viewed in these two ways.


The two nonabelian groups of order 8







The subgroup lattice of D_4



The subgroup lattice of D_4



The three abelian groups of order 8



More on subgroups

Tip

It will be *essential* to learn the subgroup lattices of our standard examples of groups.

Let's summarize the sizes of the subgroups of the groups of order 8 that we have seen.

	C_8	Q_8	$C_4 \times C_2$	D_4	C_2^3
# elts. of order 8	4	0	0	0	0
# elts. of order 4	2	6	4	2	0
# elts. of order 2	1	1	3	5	7
# elts. of order 1	1	1	1	1	1
# subgroups	4	6	8	10	16

Rule of thumb

Groups with elements of small order tend to have more subgroups than those with elements of large order.

One-step subgroup test (exercise)

A subset $H \subseteq G$ is a subgroup if and only if the following condition holds:

$$\text{If } x, y \in H, \text{ then } xy^{-1} \in H.$$

Subgroups of cyclic groups

Proposition

Every subgroup of a cyclic group is cyclic.

Proof

Let $H \leq G = \langle x \rangle$, and $|H| > 1$.

Let x^k be the smallest positive power of x in $H = \{x^k \mid k \in \mathbb{Z}\}$

We'll show that all elements of H have the form $(x^k)^m = x^{km}$ for some $m \in \mathbb{Z}$.

Take any other $x^\ell \in H$, with $\ell > 0$, and write $\ell = qk + r$, where $0 \leq r < k$.

We have $x^\ell = x^{qk+r}$, and hence

$$x^r = x^{\ell - qk} = x^\ell x^{-qk} = x^\ell (x^k)^{-q} \in H.$$

Minimality of $k > 0$ forces $r = 0$. □

Corollary


The subgroup of $G = \mathbb{Z}$ generated by a_1, \dots, a_k is $\langle \gcd(a_1, \dots, a_k) \rangle \cong \mathbb{Z}$. □

Subgroups of cyclic groups


If d divides n , then $\langle d \rangle \leq \mathbb{Z}_n$ has order n/d . Moreover, all cyclic subgroups have this form.

Corollary

The subgroups of \mathbb{Z}_n are of the form $\langle d \rangle$ for every divisor d of n . □



subgroup lattice



divisor lattice


The **order** can be read off from the divisor lattice of 24.

Cosets

Definition

Let $H \leq G$. Given $x \in G$, its **left coset** xH and **right coset** Hx are:

$$xH = \{xh \mid h \in H\}, \quad Hx = \{hx \mid h \in H\}.$$



Lagrange's theorem

Remark

For any $H \leq G$, the left cosets of H partition G into subsets of equal size (exercise).

The right cosets also partition G into subsets of equal size, but *they may be different*.

Let's compare these partitions for $H = \langle f \rangle$ in $G = D_4$.

H	r^2H	rH	r^3H
f	r^2f	rf	r^3
1	r^2	r	r^3f

H	Hr^2			
f	r^2f	fr^3	r^3	Hr^3
1	r^2	r	fr	Hr

Definition

The **index** of $H \leq G$, written $[G : H]$, is the number of distinct left (or equivalently, right) cosets of H in G .

Lagrange's theorem

If H is a subgroup of finite group G , then $|G| = [G : H] \cdot |H|$. □

The tower law

Proposition

Let G be a finite group and $K \leq H \leq G$ be a chain of subgroups. Then

$$[G : K] = [G : H][H : K].$$

Here is a “proof by picture”:

$[G : H] = \#$ of cosets of H in G

$[H : K] = \#$ of cosets of K in H

$[G : K] = \#$ of cosets of K in G

zH	z_1K	z_2K	z_3K	\cdots	z_nK
	\vdots	\vdots	\vdots	\ddots	\vdots
aH	a_1K	a_2K	a_3K	\cdots	a_nK
H	K	h_2K	h_3K	\cdots	h_nK

Proof



By Lagrange's theorem,

$$[G : H][H : K] = \frac{|G|}{|H|} \cdot \frac{|H|}{|K|} = \frac{|G|}{|K|} = [G : K]. \quad \square$$

The tower law

Another way to visualize the tower law involves subgroup lattices.

It is often helpful to label the edge from H to K in a subgroup lattice with the index $[H : K]$.





The tower law and subgroup lattices

For any two subgroups $K \leq H$ of G , the index of K in H is just the *products of the edge labels* of any path from H to K .

Equality of sets vs. equality of elements

Caveat!

An equality of cosets $xH = Hx$ as sets *does not* imply an equality of elements $xh = hx$.



rH	r	r^3	rf	r^3f
H	1	r^2	f	r^2f

r	r^3	fr	fr^3	Hr
1	r^2	f	fr^2	H

Proposition

If $[G : H] = 2$, then both left cosets of H are also right cosets.

The center of a group

Definition

The **center** of G is the set

$$Z(G) = \{z \in G \mid gz = zg, \forall g \in G\}.$$

If $z \in Z(G)$, we say that z is **central** in G .

Examples

Let's think about what elements commute with everything in the following groups:

■ $Z(D_4) = \langle r^2 \rangle = \{1, r^2\}$

■ $Z(\mathbf{Frz}_1) = \langle v \rangle = \{1, v\}$

■ $Z(D_3) = \{1\}$

■ $Z(S_4) = \{e\}$

■ $Z(Q_8) = \langle -1 \rangle = \{1, -1\}$

■ $Z(A_4) = \{e\}$

Clearly, if $H \leq Z(G)$, then $xH = Hx$ for all $x \in G$.


Proposition (exercise)

For any group G , the center $Z(G)$ is a subgroup.


Normal subgroups and normalizers

Given a subgroup H of G , it is natural to ask the following question:

How many left cosets of H are right cosets?



Partition of G by the left cosets of H



Partition of G by the right cosets of H

Definition

A subgroup H is **normal** if $gH = Hg$ for all $g \in G$. We write $H \trianglelefteq G$.

The **normalizer** of H , denoted $N_G(H)$, is the set of elements $g \in G$ such that $gH = Hg$:

$$N_G(H) = \{g \in G \mid gH = Hg\},$$

i.e., the union of left cosets that are also right cosets.

Proposition (exercise)

For any $H \leq G$,

$$H \trianglelefteq N_G(H) \leq G.$$

Conjugate subgroups

Definition

For a fixed $g \in G$, the (left) **conjugate** of H by g is

$$gHg^{-1} = \{ghg^{-1} \mid h \in H\}$$

The set of all subgroups conjugate to H is its **conjugacy class**, denoted

$$\text{cl}_G(H) = \{gHg^{-1} \mid g \in G\}.$$

Proposition (exercise)


1. gHg^{-1} is a subgroup of G ;
2. conjugation is an equivalence relation on the set of subgroups of G .

Useful remark

The following conditions are all equivalent to a subgroup $H \leq G$ being normal:

- (i) $gH = Hg$ for all $g \in G$; (“left cosets are right cosets”);
- (ii) $gHg^{-1} = H$ for all $g \in G$; (“only one conjugate subgroup”)
- (iii) $ghg^{-1} \in H$ for all $g \in G$; (“closed under conjugation”).

The alternating group A_4



Observations


- A subgroup is **normal** if its conjugacy class has size 1.
- The size of a conjugacy class tells us *how close to being normal* a subgroup is.
- Remember these subgroups:

$$|cl_{A_4}(N)| = 1 = \frac{1}{\text{Deg}_{A_4}^{\triangleleft}(N)}, \quad |cl_{A_4}(H)| = 4 = \frac{1}{\text{Deg}_{A_4}^{\triangleleft}(H)}, \quad |cl_{A_4}(K)| = 3 = \frac{1}{\text{Deg}_{A_4}^{\triangleleft}(K)}.$$

Three subgroups of A_4

The **normalizer** of each subgroup consists of the elements in the blue left cosets.


Here, take $a = (123)$, $x = (12)(34)$, $z = (13)(24)$, and $b = (234)$.



(124)	(234)	(143)	(132)
(123)	(243)	(142)	(134)
e	(12)(34)	(13)(24)	(14)(23)

$$[A_4 : N_{A_4}(N)] = 1$$


“normal”



(14)(23)	(142)	(143)
(13)(24)	(243)	(124)
(12)(34)	(134)	(234)
e	(123)	(132)

$$[A_4 : N_{A_4}(H)] = 4$$

“fully unnormal”



(124)	(234)	(143)	(132)
(123)	(243)	(142)	(134)
e	(12)(34)	(13)(24)	(14)(23)

$$[A_4 : N_{A_4}(K)] = 3$$

“moderately unnormal”

The degree of normality

Let $H \leq G$ have index $[G : H] = n < \infty$. Let's define a term that describes:

"the proportion of cosets that are blue"

Definition

Let $H \leq G$ with $[G : H] = n < \infty$. The **degree of normality** of H is

$$\text{Deg}_G^{\triangleleft}(H) := \frac{|N_G(H)|}{|G|} = \frac{1}{[G : N_G(H)]} = \frac{\# \text{ elements } x \in G \text{ for which } xH = Hx}{\# \text{ elements } x \in G}.$$

- If $\text{Deg}_G^{\triangleleft}(H) = 1$, then H is **normal**.
- If $\text{Deg}_G^{\triangleleft}(H) = \frac{1}{n}$, we'll say H is **fully unnormal**.
- If $\frac{1}{n} < \text{Deg}_G^{\triangleleft}(H) < 1$, we'll say H is **moderately unnormal**.

Big idea

The degree of normality measures *how close to being normal* a subgroup is.

A special case of the orbit-stabilizer theorem


Theorem

Let $H \leq G$ with $[G : H] = n < \infty$. Then

$$|\text{cl}_G(H)| = \frac{1}{\text{Deg}_G^{\triangleleft}(H)} = [G : N_G(H)] = \frac{\# \text{ elements } x \in G \text{ for which } xH = Hx}{\# \text{ elements } x \in G}.$$


That is, H has exactly $[G : N_G(H)]$ conjugate subgroups.

$$G = N_G(N)$$




normal

$$|\text{cl}_G(N)| = 1$$



fully unnormal

$$|\text{cl}_G(H)| = [G : H]; \text{ as large as possible}$$



moderately unnormal


$$1 < |\text{cl}_G(K)| < [G : K]$$

“Reducing” subgroup lattices

Sometimes it is convenient to collapse conjugacy classes into single nodes in the lattice.

We'll call this the **conjugacy poset** (it need not be a lattice!). Sometimes it reveals patterns in new ways.

The left-subscript denotes the size.




Conjugating normal subgroups

Proposition

If $H \leq N \trianglelefteq G$, then $xHx^{-1} \leq N$ for all $x \in G$.



Proof

Conjugating $H \leq N$ by $x \in G$ yields $xHx^{-1} \leq xNx^{-1} = N$. □



Determining the conjugacy classes by inspection

Suppose we conjugate $G = D_4$ by some element $x \in D_4$.




Remarks:

- Subgroups at a unique "lattice neighborhood," called **unicorns**, must be normal.
- all index-2 subgroups are normal.
- order-2 subgroups are normal iff they're central. (Why?)
- each nonnormal order-2 subgroup $\langle r^i f \rangle$ has a:
 - size-2 conjugacy class. (Why?)
 - index-2 normalizer, $N_{D_4}(\langle r^i f \rangle) = \langle r^i, f \rangle$.

Unicorns in the diquaternion group

Our definition of **unicorn** could be strengthened, but we want to keep things simple.


Are any of the C_4 subgroups of DQ_8 unicorns, i.e., “not like the others”?



What can we say about conjugacy classes of the subgroups of DQ_8 just from the lattice?

A mystery group of order 16

Let's repeat a previous exercise, for this lattice of an actual group. Unicorns are purple.



Every subgroup is normal, except possibly $\langle s \rangle$ and $\langle r^4 s \rangle$. (Why?)

There are two cases:

- $\langle s \rangle$ and $\langle r^4 s \rangle$ are normal $\Rightarrow s \in Z(G) \Rightarrow G$ is abelian.
- $\langle s \rangle$ and $\langle r^4 s \rangle$ are not normal $\Rightarrow \text{cl}_G(\langle s \rangle) = \{\langle s \rangle, \langle r^4 s \rangle\} \Rightarrow G$ is nonabelian.



This doesn't necessarily mean that both of these are actually possible. . .

A mystery group of order 16

It's straightforward to check that this is the subgroup lattice of

$$C_8 \times C_2 = \langle r, s \mid r^8 = s^2 = 1, srs = r \rangle.$$



Let $r = (a, 1)$ and $s = (1, b)$, and so $C_8 \times C_2 = \langle r, s \rangle = \langle (a, 1), (1, b) \rangle$.



A mystery group of order 16

However, the nonabelian case is possible as well! The following also works:


$$SA_8 = \langle r, s \mid r^8 = s^2 = 1, srs = r^5 \rangle.$$



More on conjugate subgroups


Proposition (exercise)

If $aH = bH$, then $Ha^{-1} = Hb^{-1}$, and hence $aHa^{-1} = bHb^{-1}$.




Proposition (HW)

For any $H \leq G$, the intersection of all conjugates is normal: $N := \bigcap_{x \in G} xHx^{-1} \trianglelefteq G$.








The subgroup lattice of the simple group A_5



Conjugate subgroups, visually

Consider the subgroups $A = \langle a \rangle$ and $B = \langle b \rangle$ of $G = C_4 \times C_4$.



Conjugate elements

Definition

The **conjugacy class** of an element $h \in G$ is the set

$$\text{cl}_G(h) = \{xhx^{-1} \mid x \in G\}.$$

Proposition (“class equation”)

For any finite group G ,

$$|G| = |Z(G)| + \sum |\text{cl}_G(h_i)|,$$

where the sum is taken over distinct conjugacy classes of size greater than 1.

Proof (sketch)

Immediate upon showing that:

- $|\text{cl}_G(h)| = 1$ iff $h \in Z(G)$;
- conjugacy of elements is an **equivalence relation**.

Proposition (exercise)

Every normal subgroup is the union of conjugacy classes.

Conjugate elements



Often, we can determine the conjugacy classes by inspection.

Let's look at Q_8 , all of whose subgroups are normal.

- Since $i \notin Z(Q_8) = \{\pm 1\}$, we know $|\text{cl}_{Q_8}(i)| > 1$.
- Also, $\langle i \rangle = \{\pm 1, \pm i\}$ is a union of conjugacy classes.
- Therefore $\text{cl}_{Q_8}(i) = \{\pm i\}$.

Similarly, $\text{cl}_{Q_8}(j) = \{\pm j\}$ and $\text{cl}_{Q_8}(k) = \{\pm k\}$.


1	i	j	k
-1	$-i$	$-j$	$-k$



“Conjugation preserves structure”

Revisiting frieze groups, let $h = h_0$ denote the reflection across the central axis, l_0 .

Suppose we want to reflect across a different axis, say l_{-2} .



It should be clear that all reflections (resp., rotations) of the “same parity” are conjugate.


Conjugacy classes in D_n

The dihedral group D_n is a “finite version” of the previous frieze group.

When n is even, there are two “types of reflections” of an n -gon:

1. $r^{2k}f$ is across an axis that bisects two sides
2. $r^{2k+1}f$ is across an axis that goes through two corners.

Here is a visual reason why each of these two types form a conjugacy class in D_n .



What do you think the conjugacy classes of a reflection is in D_n when n is odd?

Centralizers

Definition

The **centralizer** of $h \in G$ is the set of elements that **commute with h**

$$C_G(h) = \{x \in G \mid xh = hx\} \leq G.$$

Exercise: (i) $C_G(h)$ contains at least $\langle h \rangle$, (ii) if $xh = hx$, then $x\langle h \rangle \subseteq C_G(h)$.

Definition

Let $h \in G$ with $[G : \langle h \rangle] = n < \infty$. The **degree of centrality** of h is

$$\text{Deg}_G^C(h) := \frac{|C_G(h)|}{|G|} = \frac{1}{[G : C_G(h)]} = \frac{\# \text{ elements } x \in G \text{ for which } xh = hx}{\# \text{ elements } x \in G}.$$

- If $\text{Deg}_G^C(h) = 1$, then h is **central**.
- If $\text{Deg}_G^C(h) = \frac{1}{n}$, we'll say h is **fully uncentral**.
- If $\frac{1}{n} < \text{Deg}_G^C(h) < 1$, we'll say h is **moderately uncentral**.

Big idea

The degree of centrality measures *how close to being central* an element is.

The number of conjugate elements

The following result is analogous to an earlier one on the degree of normality and $|cl_G(H)|$.


Theorem

Let $h \in G$ with $[G : \langle h \rangle] = n < \infty$. Then

$$|cl_G(h)| = \frac{1}{\text{Deg}_G^C(h)} = [G : C_G(h)] = \frac{\# \text{ elements } x \in G \text{ for which } xh = hx}{\# \text{ elements } x \in G}.$$


That is, there are exactly $[G : C_G(h)]$ elements conjugate to h .

Both of these are special cases of the **orbit-stabilizer theorem**, about group actions.




central

$$|cl_G(h)| = 1$$



fully uncentral



$$|cl_G(h)| = [G : \langle h \rangle]; \text{ as large as possible}$$



moderately uncentral

$$1 < |cl_G(h)| < [G : \langle h \rangle]$$

An example: conjugacy classes and centralizers in Dic_6



rs	r^3s	r^5s
s	r^2s	r^4s
r^3	r^2	r^4
1	r	r^5

conjugacy classes

r^2	r^5	r^2s	r^5s
r	r^4	rs	r^4s
1	r^3	s	r^3s

$[G : C_G(r^3)] = 1$
"central"

rs	r^3s	r^5s
s	r^2s	r^4s
r	r^3	r^5
1	r^2	r^4

$[G : C_G(r^2)] = 2$
"moderately uncentral"

r^2	r^2s	r^5	r^5s
r	rs	r^4	r^4s
1	s	r^3	r^3s



$[G : C_G(s)] = 3$
"fully uncentral"

Conjugacy classes in D_6


Let's find the conjugacy classes of D_6 by inspection. The centralizers are:

- $C_{D_6}(1) = C_{D_6}(r^3) = D_6$, (order 12; index 1)
- $C_{D_6}(r^i) = \langle r \rangle$, for $i = 2, 3, 4, 5$, (order 6; index 2)
- $C_{D_6}(r^i f) = \langle r^3, r^i f \rangle = \{1, r^3, r^i f, r^{3+i} f\}$, (order 4; index 3).

This is enough information to determine the conjugacy classes!




1	r	r^2	f	$r^2 f$	$r^4 f$
r^3	r^5	r^4	$r f$	$r^3 f$	$r^5 f$






The subgroup lattice of D_6

We now can deduce the conjugacy classes of the subgroups of D_6 .



The conjugacy poset of D_6







Conjugacy classes in D_5

Since $n = 5$ is odd, all reflections in D_5 are conjugate.

Centralizers

- $C_{D_5}(1) = D_5$ (index 1),
- $C_{D_5}(r^i) = \langle r \rangle$ (index 2),
- $C_{D_5}(r^i f) = \langle r^2 f \rangle$ (index 5).



1	rf	r^3f	r	r^4
f	r^2f	r^4f	r^2	r^3

Cycle type and conjugacy in the symmetric group

We introduced **cycle type** in back in Chapter 2.

This is best seen by example. There are five cycle types in S_4 :

example element	e	(12)	(234)	(1234)	$(12)(34)$
parity	even	odd	even	odd	even
# elts	1	6	8	6	3

Definition

Two elements in S_n have the same **cycle type** if when written as a product of disjoint cycles, there are the same number of length- k cycles for each k .

Theorem

Two elements $g, h \in S_n$ are **conjugate** if and only if they have the same **cycle type**.

For example, permutations in S_5 fall into seven cycle types (conjugacy classes):

$$\text{cl}(e), \quad \text{cl}((12)), \quad \text{cl}((123)), \quad \text{cl}((1234)), \quad \text{cl}((12345)), \quad \text{cl}((12)(34)), \quad \text{cl}((12)(345)).$$

Big idea

Conjugate permutations have the same structure – they are *the same up to renumbering*.


Conjugation preserves structure in the symmetric group

The symmetric group $G = S_6$ is generated by any transposition and any n -cycle.

Consider the permutations of seating assignments around a circular table achievable by

- (23) : "people in chairs 2 and 3 may swap seats"
- (123456) : "people may cyclically rotate seats counterclockwise"

Here's how to get people in chairs 1 and 2 to swap seats:




The subgroup lattice of S_4

Exercise


Partition the subgroup lattice of S_4 into conjugacy classes by inspection alone.

Order = 24

Index = 1



The conjugacy poset of S_4



Conjugacy class size

Theorem (number of conjugate subgroups)

The **conjugacy class** of $H \leq G$ contains exactly $[G : N_G(H)]$ subgroups.

Proof (roadmap)

Construct a bijection between **left cosets** of $N_G(H)$ and **conjugate subgroups** of H :

" $xHx^{-1} = yHy^{-1}$ iff x and y are in the same left coset of $N_G(H)$."

Define $\phi: \{\text{left cosets of } N_G(H)\} \longrightarrow \{\text{conjugates of } H\}$, $\phi: xN_G(H) \longmapsto xHx^{-1}$.

Theorem (number of conjugate elements)

The **conjugacy class** of $h \in G$ contains exactly $[G : C_G(h)]$ elements.

Proof (roadmap)

Construct a bijection between **left cosets** of $C_G(h)$, and **elements** in $\text{cl}_G(h)$:

" $xhx^{-1} = yhy^{-1}$ iff x and y are in the same left coset of $C_G(h)$."

Define $\phi: \{\text{left cosets of } C_G(h)\} \longrightarrow \{\text{conjugates of } h\}$, $\phi: xC_G(h) \longmapsto xhx^{-1}$.

Quotients


Denote the set of left cosets of H in G by

$$G/H := \{xH \mid x \in G\}.$$


Key idea

The **quotient** of G by a subgroup H exists when the (left) cosets of H form a group.

This is well-defined precisely when H is **normal**.



Cluster the left cosets of N




Collapse cosets into single nodes


	N	iN	jN	kN
N	N	iN	jN	kN
iN	iN	N	kN	jN
jN	jN	kN	N	iN
kN	kN	jN	iN	N

Elements of the quotient are cosets of N


Quotients




Cluster the left cosets of $H \leq \mathbb{Z}_6$




Collapse cosets into single nodes




Elements of the quotient are cosets of H



Cluster the left cosets of $N \leq D_3$



Collapse cosets into single nodes





Elements of the quotient are cosets of N



We say that $\mathbb{Z}_6/\langle 2 \rangle \cong \mathbb{Z}_2$ and $D_3/\langle r \rangle \cong C_2$.

Quotients

Let's revisit $N = \langle (12)(34), (13)(24) \rangle$ and $H = \langle (123) \rangle$ of A_4 :




	N	aN	a^2N
N	N	aN	a^2N
aN	aN	a^2N	N
a^2N	a^2N	N	aN



When do the cosets of H form a group?

In the following: *the right coset Hg consists of the nodes at the “arrowtips”*.



Key idea

If H is **normal subgroup** of G , then the quotient group G/H exists.

If H is not normal, then following the blue arrows from H is **ambiguous**.

In other words, it **depends on our where we start within H** .

What does it mean to “multiply” two cosets?

Proposition


If $H \trianglelefteq G$, the set of left cosets G/H forms a group, with binary operation

$$aH \cdot bH := abH.$$

It's clear that G/H is closed under this operation, we just have to show that the operation is **well-defined**.

By that, we mean that it *does not depend on our choice of coset representative*:

if $a_1H = a_2H$ and $b_1H = b_2H$, then $a_1H \cdot b_1H = a_2H \cdot b_2H$.



Quotient groups, algebraically

Lemma

When $H \trianglelefteq G$, the set of cosets G/H forms a group.

Proof

To show the binary operation is, suppose $a_1H = a_2H$ and $b_1H = b_2H$. Then

$$\begin{aligned} a_1H \cdot b_1H &= a_1b_1H && \text{(by definition)} \\ &= a_1(b_2H) && (b_1H = b_2H \text{ by assumption}) \\ &= (a_1H)b_2 && (b_2H = Hb_2 \text{ since } H \trianglelefteq G) \\ &= (a_2H)b_2 && (a_1H = a_2H \text{ by assumption}) \\ &= a_2b_2H && (b_2H = Hb_2 \text{ since } H \trianglelefteq G) \\ &= a_2H \cdot b_2H && \text{(by definition)} \end{aligned}$$

Thus, the binary operation on G/H is well-defined.

We'll leave checking the group axioms as an exercise. □