

1. Let $P(t)$ be the net worth of an investment after t years, that is growing at a 5% rate. Suppose that after two years, the investment is worth \$200. Write down an *initial value problem* (the differential equation & initial condition) for P , and sketch its solution.
2. Let $T(t)$ be the temperature of a cup of water t minutes after being placed in a room where the ambient temperature is 72° .
 - (a) Write down a differential equation that T satisfies.
 - (b) Sketch the solution curve of the solution satisfying $T(0) = 100$.
 - (c) Sketch the solution curve of the solution satisfying $T(0) = 40$.
 - (d) Sketch the solution curve of the solution satisfying $T(0) = 72$.
3. Repeat the previous exercise, except let $T(t)$ be the temperature of a sheet of metal (which cools down and heats up *much* quicker than water). Qualitatively, what is the difference between the solution curves in these two problems? Which value of k is bigger and why?
4. Sketch the slope field of the ODE $y' = t - 2y$ using the isocline method for $y' = c$, for $c = 0, \pm 1, \pm 2, \pm 3$. Sketch the particular solutions that satisfy $y(0) = 1$ and $y(2) = 2$.
5. Sketch the slope field of the ODE $y' = -t$ using the isocline method for $c = 0, \pm \frac{1}{2}, \pm 1, \pm 2$, and sketch the particular solution that satisfies $y(0) = 1$.
6. Let $y' = f(y, t)$ be an ODE. Explain why two solution curves in its slope field can never cross.
7. Sketch the steady-state (constant) solutions of $y' = 6 + y - y^2$ in the ty -plane. These solutions divide the plane into regions. Sketch at least one solution curve in each of these region.
8. Consider the differential equation $y' = y(4 - y)$.
 - (a) Show that $y(t) = 4/(1 + Ce^{-4t})$ is a solution for any value of C by plugging it into the ODE. This family of solutions is called a *general solution* to the differential equation.
 - (b) Sketch the solutions for $C = 1, 2, \dots, 5$. (Hint: This ODE is autonomous).
 - (c) What are the steady-state (constant) solutions?
 - (d) The general solution may fail to produce all solutions of a differential equation. Find a solution that is not given by any value of C . (Hint: Look at part (c)).
 - (e) Describe a physical situation that this differential equation could model, and justify your reasons. (Hint: Consider population growth).
9. Consider the initial value problem $y' = t + y$, $y(0) = 1$.
 - (a) When computing a solution by hand using Euler's method, it is beneficial to arrange your work in a table, as shown below where the first step is computed. Continue with Euler's method using step-size $h = 0.1$ and complete all missing entries of the table.

k	t_k	y_k	$f(t_k, y_k) = t_k + y_k$	h	$f(t_k, y_k) \cdot h$
0	0.0	1.0	1.0	0.1	0.1
1	0.1	1.1			
2	0.2				
3	0.3				
4	0.4				
5	0.5				

- (b) The general solution of $y' = t + y$ is $y(t) = Ce^t - t - 1$. Using this, compute the actual value of $y(0.5)$.
10. Consider the initial value problem $y' = (1 + t)y$, $y(0) = -1$.
- (a) Use Euler's method to approximate $y(1)$, for step-size $h = 0.2$, and then for $h = 0.1$. Arrange your results in the tabular form as in the previous exercise.
- (b) Compute the actual value of $y(1)$ by solving the initial value problem $y' = (1 + t)y$, $y(0) = -1$ and plugging in $t = 1$.