MthSc 208: Differential Equations (Summer II, 2010) In-class Worksheet 7c: The 2D Heat Equation

NAME:

We will solve for the function u(x, y, t) defined for $0 \le x, y \le \pi$ and $t \ge 0$ which satisfies the following initial value problem of the heat equation:

$$u_t = c^2(u_{xx} + u_{yy})$$

$$u(0, y, t) = u(\pi, y, t) = u(x, 0, t) = u(x, \pi, t) = 0,$$

$$u(x, y, 0) = 2\sin x \sin 2y + 3\sin 4x \sin 5y.$$

(a) Carefully describe (and sketch) a physical situation that this models.

(b) Assume that u(x, y, t) = f(x, y)g(t). Compute u_{xx} , u_{yy} , and u_t , find boundary conditions for f(x, y).

(c) Plug u = fg back into the PDE and separate variables by dividing both sides of the equation by $c^2 fg$. Set this equal to a constant λ , and write down two equations: an ODE for g(t), and a PDE f(x,y) (called the *Helmholtz equation*), with four boundary conditions.

(d) Solve the ODE for g(t).

(e) To solve the PDE for f, assume that f(x,y) = X(x)Y(y). Plug this back in and separate variables. [For consistency, put the X''/X term on one side of the equation, and set equal to a constant μ .]

(f) Write down two ODEs – one for X(x) and one for Y(y), and include boundary conditions for both. Hint: It is easier notationally if you introduce a new constant, $\nu := \lambda - \mu$.

(g) Solve the ODEs for X(x) and Y(y), and determine μ and ν (and hence λ). You should get a λ for each choice of positive integers $n, m \in \mathbb{N}$, call it λ_{nm} .

(h) For each $n, m \in \mathbb{N}$, we have a solution $u_{nm}(x, y, t) = f_{nm}(x, y)g_{nm}(t)$. Write down this solution.

(i) Find the general solution of the PDE. It will be a doubly infinite sum (superposition) of solutions: $\sum_{n,m\in\mathbb{N}}u_{nm}(x,y,t).$

(g) Find the particular solution to the initial value problem by using the initial condition.

(h) What is the long-term behavior of the system? Give a mathematical, and physical, justification.