MthSc 208: Differential Equations (Summer II, 2012) In-class Worksheet 6c: Parseval's Identity

NAME:

Recall that Parseval's identity says that

$$\frac{1}{\pi} \int_{-\pi}^{\pi} (f(x))^2 dx = \frac{1}{2} a_0^2 + \sum_{n=1}^{\infty} (a_n^2 + b_n^2).$$

We will use this to compute $\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \cdots$.

- 1. Let f(x) = x on $[-\pi, \pi]$ and extend f(x) to be 2π -periodic. Write f(x) as a Fourier series. (See Example 2 on pages 5-6 of the lecture notes.)
- 2. Compute $\frac{1}{\pi} \int_{-\pi}^{\pi} (f(x))^2 dx$. (The left-hand side of Parseval's identity.)

3. Compute $\frac{1}{2}a_0^2 + \sum_{n=1}^{\infty} (a_n^2 + b_n^2)$. (The right-hand side of Parseval's identity.)

4. Equate your answers to the previous two parts and solve for $\sum_{n=1}^{\infty} \frac{1}{n^2}$.