

1. For each of the first-order differential equations, decide whether it is linear or nonlinear. If the equation is linear, state whether it is homogeneous or inhomogeneous.

(a) $y' = ky$

(b) $y' = k(72 - y)$

(c) $y' = y(4 - y)$

(d) $y' = t + y$

(e) $3y' + 5y = 3 \cos 2t$

(f) $3y' + 5y = 3 \cos 2y$

(g) $y' = 4t^2y - \sin t$

(h) $y' = 4ty^2 - \sin t$

2. Use the integrating factor method to find the general solution of the following differential equations.

(a) $2y' - 3y = 5$

(b) $y' + 2ty = 5t$

(c) $ty' = 4y + t^4$

3. Use the variation of parameters method to find the general solution of the following differential equation. Then find the particular solution satisfying the given initial condition.

(a) $y' - 3y = 4, \quad y(0) = 2$

(b) $y' + y = e^t, \quad y(0) = 1$

(c) $y' + 2ty = 2t^3, \quad y(0) = -1.$

4. Solve the differential equation $y' = 2y + 4$ four different ways:

(a) Undetermined coefficients (that is, $y(t) = y_h(t) + y_p(t)$)

(b) Integrating factor

(c) Variation of parameters

(d) Separation of variables.

5. Solve the following ODEs using the method of undetermined coefficients: That is, write $y(t) = y_h(t) + y_p(t)$, then solve the homogeneous equation, then guess a particular solution, and add those together to get the general solution.

(a) $y' - 2y = 0$

(b) $y' - 2y = 10$

(c) $y' - 2y = t$

(d) $y' - 2y = t^2 + 1$

(e) $y' - 2y = 4e^{3t}$

(f) $y' - 2y = 5 \sin 3t$

6. A tank contains 100 gal of pure water. A salt solution with concentration 3 lb/gal enters the tank at a rate of 2 gal/min. Solution drains from the tank at a rate of 2 gal/min. Sketch a picture of this situation, then *without doing any math*, determine the eventual concentration of the salt solution in the tank (i.e., the steady-state solution).
7. A tank contains 100 gal of pure water. At time zero, a sugar-water solution containing 0.2 lb of sugar per gal enters the tank at a rate of 3 gal per minute. Simultaneously, a drain is opened at the bottom of the tank allowing the sugar solution to leave the tank at 3 gal per minute. Assume that the solution in the tank is kept perfectly mixed at all time.
- (a) What will be the sugar content in the tank after 20 minutes?
- (b) How long will it take the sugar content in the tank to reach 15 lb?
- (c) What will be the eventual sugar content in the tank?
8. A tank contains 500 gal of a salt-water solution containing 0.05 lb of salt per gallon of water. Pure water is poured into the tank and a drain at the bottom of the tank is adjusted so as to keep the volume of solution in the tank constant. At what rate (gal/min) should the water be poured into the tank to lower the salt concentration to 0.01 lb/gal of water in under one hour (i.e., $t = 60$ minutes)?
9. A tank initially contains 100 gal of a salt-water solution containing 0.05 lb of salt for each gallon of water. At time zero, pure water is poured into the tank at a rate of 2 gal per minute. Simultaneously, a drain is opened at the bottom of the tank that allows the salt-water solution to leave at a rate of 3 gal per minute. What will be the salt content in the tank when precisely 50 gal of salt solution remain.
10. A murder victim is discovered at midnight at the temperature of the body is recorded at 31°C , and it was discovered that the proportionality constant in Newton's law was $k = \ln(5/4)$. Assume that at midnight the surrounding air temperature $A(t)$ is 0°C , and is falling at a constant rate of 1°C per hour. At what time did the victim die? (Set $T(t) = 37$ and solve for t – use a computer or calculator for this part.) [*Hint*: Letting $t = 0$ represent midnight will simplify your calculations.]