

## Chapter 10: The Sylow Theorems

Matthew Macauley

Department of Mathematical Sciences  
Clemson University  
<http://www.math.clemson.edu/~macaule/>

Math 4120, Summer I 2014

## Overview

This chapter about one question:

What groups are there?

In Chapter 5, we saw five families of groups: cyclic, dihedral, abelian, symmetric, alternating.

In Chapter 8, we classified all (finitely generated) *abelian* groups.

But what *other* groups are there, and what do they look like? For example, for a fixed order  $|G|$ , we may ask the following questions about  $G$ :

1. How big are its subgroups?
2. How are those subgroups related?
3. How many subgroups are there?
4. Are any of them normal?

There is no one general method to answer this for any given order.

However, the **Sylow Theorems**, developed by Norwegian mathematician Peter Sylow (1832–1918), are powerful tools that help us attack this question.

# The Sylow Theorems

## Definition

A  **$p$ -group** is a group whose order is a power of a prime  $p$ . A  $p$ -group that is a subgroup of a group  $G$  is a  **$p$ -subgroup** of  $G$ .

## Notational convention

Throughout,  $G$  will be a group of order  $|G| = p^n \cdot m$ , with  $p \nmid m$ . That is,  $p^n$  is the *highest power of  $p$  dividing  $|G|$* .


There are three **Sylow theorems**, and loosely speaking, they describe the following about a group's  $p$ -subgroups:

1. **Existence:** In every group,  $p$ -subgroups of all possible sizes exist.
2. **Relationship:** All maximal  $p$ -subgroups are conjugate.
3. **Number:** There are strong restrictions on the number of  $p$ -subgroups a group can have.

Together, these place strong restrictions on the structure of a group  $G$  with a fixed order.

## Our unknown group of order 200

Throughout this chapter, we will have a running example, a “mystery group”  $M$  of order 200.



Using *only* the fact that  $|M| = 200$ , we will uncover as much about the structure of  $M$  as we can.

We actually already know a little bit. Recall Cauchy's theorem:

### Cauchy's theorem


If  $p$  is a prime number dividing  $|G|$ , then  $G$  has an element  $g$  of order  $p$ .

## Our mystery group of order 200

Since our mystery group  $M$  has order  $|M| = 2^3 \cdot 5^2 = 200$ , Cauchy's theorem tells us that:

- $M$  has an element  $a$  of order 2;
- $M$  has an element  $b$  of order 5;

Also, by Lagrange's theorem,  $\langle a \rangle \cap \langle b \rangle = \{e\}$ .



## $p$ -groups

Before we introduce the Sylow theorems, we need to better understand  $p$ -groups.

Recall that a  $p$ -group is any group of order  $p^n$ . For example,  $C_1$ ,  $C_4$ ,  $V_4$ ,  $D_4$  and  $Q_4$  are all 2-groups.

### $p$ -group Lemma


If a  $p$ -group  $G$  acts on a set  $S$  via  $\phi: G \rightarrow \text{Perm}(S)$ , then

$$|\text{Fix}(\phi)| \equiv_p |S|.$$

### Proof (sketch)

Suppose  $|G| = p^n$ .

By the Orbit-Stabilizer theorem, the only possible orbit sizes are  $1, p, p^2, \dots, p^n$ .



## $p$ -groups

### Normalizer lemma, Part 1

If  $H$  is a  $p$ -subgroup of  $G$ , then

$$[N_G(H) : H] \equiv_p [G : H].$$

### Proof

Let  $S = G/H = \{Hx \mid x \in G\}$ . The group  $H$  acts on  $S$  by **right-multiplication**, via  $\phi: H \rightarrow \text{Perm}(S)$ , where

$\phi(h)$  = the permutation sending each  $Hx$  to  $Hxh$ .

The **fixed points** of  $\phi$  are the cosets  $Hx$  in the **normalizer**  $N_G(H)$ :

$$\begin{aligned} Hxh = Hx, \quad \forall h \in H &\iff Hxhx^{-1} = H, \quad \forall h \in H \\ &\iff xhx^{-1} \in H, \quad \forall h \in H \\ &\iff x \in N_G(H). \end{aligned}$$

Therefore,  $|\text{Fix}(\phi)| = [N_G(H) : H]$ , and  $|S| = [G : H]$ . By our  $p$ -group Lemma,


$$|\text{Fix}(\phi)| \equiv_p |S| \implies [N_G(H) : H] \equiv_p [G : H].$$

□

## $p$ -groups

Here is a picture of the action of the  $p$ -subgroup  $H$  on the set  $S = G/H$ , from the proof of the Normalizer Lemma.

$S = G/H =$  set of right cosets of  $H$  in  $G$



The fixed points are precisely the cosets in  $N_G(H)$

Orbits of size  $> 1$  are of various sizes dividing  $|H|$ , but all lie outside  $N_G(H)$




## $p$ -subgroups

The following result will be useful in proving the first Sylow theorem.

### The Normalizer lemma, Part 2

Suppose  $|G| = p^n m$ , and  $H \leq G$  with  $|H| = p^i < p^n$ . Then  $H \subsetneq N_G(H)$ , and the index  $[N_G(H) : H]$  is a multiple of  $p$ .



Conclusions:

- $H = N_G(H)$  is impossible!
- $p^{i+1}$  divides  $|N_G(H)|$ .

## Proof of the normalizer lemma

### The Normalizer lemma, Part 2

Suppose  $|G| = p^n m$ , and  $H \leq G$  with  $|H| = p^i < p^n$ . Then  $H \not\leq N_G(H)$ , and the index  $[N_G(H) : H]$  is a multiple of  $p$ .

### Proof

Since  $H \triangleleft N_G(H)$ , we can create the quotient map

$$q: N_G(H) \longrightarrow N_G(H)/H, \quad q: g \longmapsto gH.$$

The size of the quotient group is  $[N_G(H) : H]$ , the number of cosets of  $H$  in  $N_G(H)$ .

By The Normalizer lemma Part 1,  $[N_G(H) : H] \equiv_p [G : H]$ . By Lagrange's theorem,

$$[N_G(H) : H] \equiv_p [G : H] = \frac{|G|}{|H|} = \frac{p^n m}{p^i} = p^{n-i} m \equiv_p 0.$$

Therefore,  $[N_G(H) : H]$  is a multiple of  $p$ , so  $N_G(H)$  must be strictly larger than  $H$ .  $\square$

# The 1<sup>st</sup> Sylow Theorem: Existence of $p$ -subgroups

## First Sylow Theorem

$G$  has a subgroup of order  $p^k$ , for each  $p^k$  dividing  $|G|$ . Also, every  $p$ -subgroup with fewer than  $p^n$  elements sits inside one of the larger  $p$ -subgroups.

The First Sylow Theorem is in a sense, a generalization of Cauchy's theorem. Here is a comparison:

Cauchy's Theorem	First Sylow Theorem
<i>If <math>p</math> divides <math> G </math>, then ...</i> There is a subgroup of order $p$ which is cyclic and has no non-trivial proper subgroups. $G$ contains an element of order $p$	<i>If <math>p^k</math> divides <math> G </math>, then ...</i> There is a subgroup of order $p^k$ which has subgroups of order $1, p, p^2 \dots p^k$ . $G$ might not contain an element of order $p^k$ .

# The 1<sup>st</sup> Sylow Theorem: Existence of $p$ -subgroups

## Proof

The trivial subgroup  $\{e\}$  has order  $p^0 = 1$ .

Big idea: Suppose we're given a subgroup  $H < G$  of order  $p^i < p^n$ . We will construct a subgroup  $H'$  of order  $p^{i+1}$ .

By the normalizer lemma,  $H \leq N_G(H)$ , and the order of the quotient group  $N_G(H)/H$  is a multiple of  $p$ .

By Cauchy's Theorem,  $N_G(H)/H$  contains an element (a coset!) of order  $p$ . Call this element  $aH$ . Note that  $\langle aH \rangle$  is cyclic of order  $p$ .


*Claim*: The **preimage** of  $\langle aH \rangle$  under the quotient  $q: N_G(H) \rightarrow N_G(H)/H$  is the subgroup  $H'$  we seek.

The preimages  $q^{-1}(H), q^{-1}(aH), q^{-1}(a^2H), \dots, q^{-1}(a^{p-1}H)$  are all distinct cosets of  $H$  in  $N_G(H)$ , each of size  $p^i$ .

Thus, the preimage  $H' = q^{-1}(\langle aH \rangle)$  contains  $p \cdot |H| = p^{i+1}$  elements. □

# The 1<sup>st</sup> Sylow Theorem: Existence of $p$ -subgroups

Here is a picture of how we found the group  $H' = q^{-1}(\langle aH \rangle)$ .




Since  $|H| = p^i$ , the subgroup  $H' = \bigcup_{k=0}^{p-1} a^k H$  contains  $p \cdot |H| = p^{i+1}$  elements.

## Our unknown group of order 200

We now know a little bit more about the structure of our mystery group of order  $|M| = 2^3 \cdot 5^2$ :

- $M$  has a 2-subgroup  $P_2$  of order  $2^3 = 8$ ;
- $M$  has a 5-subgroup  $P_5$  of order  $25 = 5^2$ ;
- Each of these subgroups contains a nested chain of  $p$ -subgroups, down to the trivial group,  $\{e\}$ .



## The 2<sup>nd</sup> Sylow Theorem: Relationship among $p$ -subgroups

### Definition

A subgroup  $H < G$  of order  $p^n$ , where  $|G| = p^n \cdot m$  with  $p \nmid m$  is called a **Sylow  $p$ -subgroup** of  $G$ . Let  $\text{Syl}_p(G)$  denote the set of Sylow  $p$ -subgroups of  $G$ .

### Second Sylow Theorem

Any two Sylow  $p$ -subgroups are conjugate (and hence isomorphic).

### Proof

Let  $H < G$  be any Sylow  $p$ -subgroup of  $G$ , and let  $S = G/H = \{gH \mid g \in G\}$ , the set of right cosets of  $H$ .

Pick *any other* Sylow  $p$ -subgroup  $K$  of  $G$ . (If there is none, the result is trivial.)

The group  $K$  acts on  $S$  by **right-multiplication**, via  $\phi: K \rightarrow \text{Perm}(S)$ , where

$$\phi(k) = \text{the permutation sending each } Hg \text{ to } Hgk.$$

## The 2<sup>nd</sup> Sylow Theorem: All Sylow $p$ -subgroups are conjugate

### Proof

A **fixed point** of  $\phi$  is a coset  $Hg \in S$  such that

$$\begin{aligned} Hgk = Hg, \quad \forall k \in K &\iff Hgkg^{-1} = H, \quad \forall k \in K \\ &\iff gkg^{-1} \in H, \quad \forall k \in K \\ &\iff gKg^{-1} \subset H \\ &\iff gKg^{-1} = H. \end{aligned}$$

Thus, if  $\phi$  has a fixed point  $gH$ , then  $H$  and  $K$  are conjugate by  $g$ , and we're done!

All we need to do is show that  $|\text{Fix}(\phi)| \not\equiv_p 0$ .

By the  $p$ -group Lemma,  $|\text{Fix}(\phi)| \equiv_p |S|$ . Recall that  $|S| = [G, H]$ .

Since  $H$  is a Sylow  $p$ -subgroup,  $|H| = p^n$ . By Lagrange's Theorem,

$$|S| = [G : H] = \frac{|G|}{|H|} = \frac{p^n m}{p^n} = m, \quad p \nmid m.$$


Therefore,  $|\text{Fix}(\phi)| \equiv_p m \not\equiv_p 0$ . □



## Our unknown group of order 200

We now know even more about the structure of our mystery group  $M$ , of order  $|M| = 2^3 \cdot 5^2$ :

- If  $M$  has any other Sylow 2-subgroup, it is isomorphic to  $P_2$ ;
- If  $M$  has any other Sylow 5-subgroup, it is isomorphic to  $P_5$ .



## The 3<sup>rd</sup> Sylow Theorem: Number of $p$ -subgroups

### Third Sylow Theorem

Let  $n_p$  be the number of Sylow  $p$ -subgroups of  $G$ . Then

$$n_p \text{ divides } |G| \quad \text{and} \quad n_p \equiv_p 1.$$

(Note that together, these imply that  $n_p \mid m$ , where  $|G| = p^n \cdot m$ .)

### Proof

The group  $G$  acts on  $S = \text{Syl}_p(G)$  by **conjugation**, via  $\phi: G \rightarrow \text{Perm}(S)$ , where

$$\phi(g) = \text{the permutation sending each } H \text{ to } g^{-1}Hg.$$

By the Second Sylow Theorem, all Sylow  $p$ -subgroups are conjugate! Thus there is **only one orbit**,  $\text{Orb}(H)$ , of size  $n_p = |S|$ .

By the Orbit-Stabilizer Theorem,

$$\underbrace{|\text{Orb}(H)|}_{=n_p} \cdot |\text{Stab}(H)| = |G| \quad \implies \quad n_p \text{ divides } |G|.$$

## The 3<sup>rd</sup> Sylow Theorem: Number of $p$ -subgroups

### Proof (cont.)

Now, pick any  $H \in \text{Syl}_p(G) = S$ . The group  $H$  acts on  $S$  by **conjugation**, via  $\theta: H \rightarrow \text{Perm}(S)$ , where

$$\theta(h) = \text{the permutation sending each } K \text{ to } h^{-1}Kh.$$

Let  $K \in \text{Fix}(\theta)$ . Then  $K \leq G$  is a Sylow  $p$ -subgroup satisfying

$$h^{-1}Kh = K, \quad \forall h \in H \quad \Longleftrightarrow \quad H \leq N_G(K) \leq G.$$

We know that:

- $H$  and  $K$  are Sylow  $p$ -subgroups of  $G$ , **but also of  $N_G(K)$** .
- Thus,  $H$  and  $K$  are conjugate in  $N_G(K)$ . (2nd Sylow Thm.)
- $K \triangleleft N_G(K)$ , thus the only conjugate of  $K$  in  $N_G(K)$  is itself.

Thus,  $K = H$ . That is,  $\text{Fix}(\theta) = \{H\}$  contains only 1 element.

By the  $p$ -group Lemma,  $n_p := |S| \equiv_p |\text{Fix}(\theta)| = 1$ .

□

# Summary of the proofs of the Sylow Theorems

For the 1st Sylow Theorem, we started with  $H = \{e\}$ , and inductively created larger subgroups of size  $p, p^2, \dots, p^n$ .

For the 2<sup>nd</sup> and 3<sup>rd</sup> Sylow Theorems, we used a clever group action and then applied one or both of the following:

- (i) *Orbit-Stabilizer Theorem*. If  $G$  acts on  $S$ , then  $|\text{Orb}(s)| \cdot |\text{Stab}(s)| = |G|$ .
- (ii)  *$p$ -group Lemma*. If a  $p$ -group acts on  $S$ , then  $|S| \equiv_p |\text{Fix}(\phi)|$ .


To summarize, we used:

- S2 The action of  $K \in \text{Syl}_p(G)$  on  $S = G/H$  by **right multiplication** for some other  $H \in \text{Syl}_p(G)$ .
- S3a The action of  $G$  on  $S = \text{Syl}_p(G)$ , by **conjugation**.
- S3b The action of  $H \in \text{Syl}_p(G)$  on  $S = \text{Syl}_p(G)$ , by **conjugation**.


## Our unknown group of order 200

We now know a little bit more about the structure of our mystery group  $M$ , of order  $|M| = 2^3 \cdot 5^2 = 200$ :

- $n_5 \mid 8$ , thus  $n_5 \in \{1, 2, 4, 8\}$ . But  $n_5 \equiv_5 1$ , so  $n_5 = 1$ .
- $n_2 \mid 25$  and is odd. Thus  $n_2 \in \{1, 5, 25\}$ .
- We conclude that  $M$  has a unique (and hence normal) **Sylow 5-subgroup**  $P_5$  (of order  $5^2 = 25$ ), and either 1, 5, or 25 **Sylow 2-subgroups** (of order  $2^3 = 8$ ).



## Our unknown group of order 200



Suppose  $M$  has a subgroup isomorphic to  $D_4$ .

This would be a Sylow 2-subgroup. Since all of them are conjugate,  $M$  *cannot* contain a subgroup isomorphic to  $Q_4$ ,  $C_4 \times C_2$ , or  $C_8$ !

In particular,  $M$  cannot even contain an element of order 8. (Why?)

# Simple groups

## Definition

A group  $G$  is **simple** if its only normal subgroups are  $G$  and  $\langle e \rangle$ .

Since all Sylow  $p$ -subgroups are **conjugate**, the following result is straightforward:

## Proposition (HW)

A Sylow  $p$ -subgroup is **normal** in  $G$  if and only if it is the **unique** Sylow  $p$ -subgroup (that is, if  $n_p = 1$ ).

The Sylow theorems are very useful for establishing statements like:

*There are no simple groups of order  $k$  (for some  $k$ ).*

To do this, we usually just need to show that  $n_p = 1$  for some  $p$  dividing  $|G|$ .

Since we established  $n_5 = 1$  for our running example of a group of size  $|M| = 200$ , there are no simple groups of order 200.

## Simple groups: an easy example

### Tip

When trying to show that  $n_p = 1$ , it's usually more helpful to analyze the largest primes first.

### Proposition

There are no simple groups of order 84.

### Proof

Since  $|G| = 84 = 2^2 \cdot 3 \cdot 7$ , the Third Sylow Theorem tells us:

- $n_7$  divides  $2^2 \cdot 3 = 12$  (so  $n_7 \in \{1, 2, 3, 4, 6, 12\}$ )
- $n_7 \equiv_7 1$ .

The only possibility is that  $n_7 = 1$ , so the Sylow 7-subgroup must be normal.  $\square$

Observe why it is beneficial to use the largest prime first:

- $n_3$  divides  $2^2 \cdot 7 = 28$  and  $n_3 \equiv_3 1$ . Thus  $n_3 \in \{1, 2, 4, 7, 14, 28\}$ .
- $n_2$  divides  $3 \cdot 7 = 21$  and  $n_2 \equiv_2 1$ . Thus  $n_2 \in \{1, 3, 7, 21\}$ .



## Simple groups: a harder example

### Proposition

There are no simple groups of order 351.

### Proof

Since  $|G| = 351 = 3^3 \cdot 13$ , the Third Sylow Theorem tells us:

- $n_{13}$  divides  $3^3 = 27$  (so  $n_{13} \in \{1, 3, 9, 27\}$ )
- $n_{13} \equiv_{13} 1$ .

The only possibilities are  $n_{13} = 1$  or 27.

A Sylow 13-subgroup  $P$  has order 13, and a Sylow 3-subgroup  $Q$  has order  $3^3 = 27$ . Therefore,  $P \cap Q = \{e\}$ .

**Suppose  $n_{13} = 27$ .** Every Sylow 13-subgroup contains 12 non-identity elements, and so  $G$  must contain  $27 \cdot 12 = 324$  elements of order 13.

This leaves  $351 - 324 = 27$  elements in  $G$  not of order 13. Thus,  $G$  contains only one Sylow 3-subgroup (i.e.,  $n_3 = 1$ ) and so  $G$  cannot be simple.  $\square$

## Simple groups: the hardest example

### Proposition

If  $H \leq G$  and  $|G|$  does not divide  $[G : H]!$ , then  $G$  cannot be simple.

### Proof

Let  $G$  act on the **right cosets** of  $H$  (i.e.,  $S = G/H$ ) by **right-multiplication**:

$$\phi: G \longrightarrow \text{Perm}(S) \cong S_n, \quad \phi(g) = \text{the permutation that sends each } Hx \text{ to } Hxg.$$

Recall that the **kernel** of  $\phi$  is the intersection of all conjugate subgroups of  $H$ :

$$\text{Ker } \phi = \bigcap_{x \in G} x^{-1} H x.$$

Notice that  $\langle e \rangle \leq \text{Ker } \phi \leq H \leq G$ , and **Ker**  $\phi \triangleleft G$ .

If  $\text{Ker } \phi = \langle e \rangle$  then  $\phi: G \hookrightarrow S_n$  is an **embedding**. But this is *impossible* because  $|G|$  does not divide  $|S_n| = [G : H]!$ . □

### Corollary

There are no simple groups of order 24.

## Theorem (classification of finite simple groups)

Every finite simple group is isomorphic to one of the following groups:

- A cyclic group  $\mathbb{Z}_p$ , with  $p$  prime;
- An alternating group  $A_n$ , with  $n \geq 5$ ;
- A Lie-type Chevalley group:  $\text{PSL}(n, q)$ ,  $\text{PSU}(n, q)$ ,  $\text{PsP}(2n, p)$ , and  $P\Omega^\epsilon(n, q)$ ;
- A Lie-type group (twisted Chevalley group or the Tits group):  $D_4(q)$ ,  $E_6(q)$ ,  $E_7(q)$ ,  $E_8(q)$ ,  $F_4(q)$ ,  ${}^2F_4(2^n)'$ ,  $G_2(q)$ ,  ${}^2G_2(3^n)$ ,  ${}^2B(2^n)$ ;
- One of 26 exceptional “sporadic groups.”

The two largest sporadic groups are the:

- “baby monster group”  $B$ , which has order

$$|B| = 2^{41} \cdot 3^{13} \cdot 5^6 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 31 \cdot 47 \approx 4.15 \times 10^{33};$$










- “monster group”  $M$ , which has order

$$|M| = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \approx 8.08 \times 10^{53}.$$

The proof of this classification theorem is spread across  $\approx 15,000$  pages in  $\approx 500$  journal articles by over 100 authors, published between 1955 and 2004.

# The Periodic Table Of Finite Simple Groups

[illegible]

-  Alternating Groups
-  Classical Chevalley Groups
-  Chevalley Groups
-  Classical Steinberg Groups
-  Steinberg Groups
-  Suzuki Groups
-  Ree Groups and Tits Group\*
-  Sporadic Groups
-  Cyclic Groups

Alternatives <sup>b</sup>	Symbol
---------------------------	--------

0-1-1

dic groups and families, alternate names are left are either names by which they known. For specific nonsynaptic groups used to indicate homophony. All such names appear on the table except the families, *no. 1* (1971).

<sup>†</sup>For sporadic groups and families, alternate names in the upper left are other names by which they may be known. For specific nonsporadic groups these are used to indicate isomorphisms. All such isomorphisms appear on the table except the families  $(\mathbf{A}_n, \mathbf{A}_n^{\text{alt}}, \mathbf{A}_n^{\text{sym}}, \mathbf{A}_n^{\text{alt}})$ .

Copyright © 2004 John Wiley & Sons, Ltd.

<sup>†</sup>These simple groups are determined by their order with the following exceptions:

- (i)  $\text{Sp}(q)$  and  $\text{O}(q)$  for  $q$  odd,  $n \geq 2$
- (ii)  $A_4$ ,  $B_4$  and  $A_5$  of order 20, 120.

$M_{11}$	$M_{12}$	$M_{22}$	$M_{23}$	$M_{24}$	$f(V), f(11)$ $f_1$	$H$ $f_2$	$H/M$ $f_3$	$J_4$	$HS$	$McL$	$He$	$Rn$
7930	95040	443 520	10 280 960	244 823 040	375 560	604 800	20 232 960	86 775 073 046 877 362 580	44 352 000	898 128 000	6 500 367 200	1 650 26 144 000

$S_z$	$O'NS, O-S$ $O'N$	-3	$Co_3$	-2	$Co_2$	-1	$Co_1$	$F_n D$ $HN$	$Ly_5$ $Ly$	$F_n E$ $Th$	$M(22)$ $F_{122}$	$M(23)$ $F_{123}$	$F_{124}, M(24)^a$ $F'_{124}$	$F_5$ $B$	$F_n M_1$ $M$
448 545 687 400	860 151 585 920	495 766 656 800	42 305 421 312 000	4 157 776 064 543 560 000	272 000 343 560 000	213 000 240 000 000	52 765 179 90 787 000	29 765 179 90 787 000	52 765 179 90 787 000	44 561 754 654 400	4 089 026 473 1 255 205 799 590	1 255 205 799 590 461 721 282 560	461 721 282 560 461 721 282 560	461 721 282 560 461 721 282 560	461 721 282 560 461 721 282 560

# Finite Simple Group (of Order Two), by The Klein Four™

## Musical Fruitcake

[View More by This Artist](#)

### Klein Four

Open iTunes to preview, buy, and download music.



[View in iTunes](#)

**\$9.99**

Genres: [Pop](#), [Music](#)

Released: Dec 05, 2005

© 2005 Klein Four

### Customer Ratings

★★★★★ 13 Ratings

	Name	Artist	Time	Price	
1	Power of One	<a href="#">Klein Four</a>	5:16	\$0.99	<a href="#">View In iTunes ▶</a>
2	Finite Simple Group (of Order Two)	<a href="#">Klein Four</a>	3:00	\$0.99	<a href="#">View In iTunes ▶</a>
3	Three-Body Problem	<a href="#">Klein Four</a>	3:17	\$0.99	<a href="#">View In iTunes ▶</a>
4	Just the Four of Us	<a href="#">Klein Four</a>	4:19	\$0.99	<a href="#">View In iTunes ▶</a>
5	Lemma	<a href="#">Klein Four</a>	3:43	\$0.99	<a href="#">View In iTunes ▶</a>
6	Calculating	<a href="#">Klein Four</a>	4:09	\$0.99	<a href="#">View In iTunes ▶</a>
7	XX Potential	<a href="#">Klein Four</a>	3:42	\$0.99	<a href="#">View In iTunes ▶</a>
8	Confuse Me	<a href="#">Klein Four</a>	3:41	\$0.99	<a href="#">View In iTunes ▶</a>
9	Universal	<a href="#">Klein Four</a>	4:13	\$0.99	<a href="#">View In iTunes ▶</a>
10	Contradiction	<a href="#">Klein Four</a>	3:48	\$0.99	<a href="#">View In iTunes ▶</a>
11	Mathematics Paradise	<a href="#">Klein Four</a>	3:51	\$0.99	<a href="#">View In iTunes ▶</a>
12	Stefanie (The Ballad of Galois)	<a href="#">Klein Four</a>	4:51	\$0.99	<a href="#">View In iTunes ▶</a>
13	Musical Fruitcake (Pass it Around)	<a href="#">Klein Four</a>	2:50	\$0.99	<a href="#">View In iTunes ▶</a>
14	Abandon Soap	<a href="#">Klein Four</a>	2:17	\$0.99	<a href="#">View In iTunes ▶</a>

14 Songs