### Lecture 3.6: Quotient, remainder, ceiling and floor

Matthew Macauley

Department of Mathematical Sciences Clemson University http://www.math.clemson.edu/~macaule/

Math 4190, Discrete Mathematical Structures

### Division and remainder

#### Theorem

Given any  $n \in \mathbb{Z}$  and  $d \in \mathbb{Z}^+$ , there exists unique integers q, r such that

 $n = dq + r, \qquad \qquad 0 \le r < d.$ 

We call  $q := n \operatorname{div} d$  and  $r := n \operatorname{mod} d$  the quotient and remainder, respectively.

#### Examples

- 1. Compute 365 div 7 and 365 mod 7.
- 2. Suppose  $n \mod 11 = 6$ . Compute  $4n \mod 11$ .
- 3. Given  $n \in \mathbb{Z}$ , compute  $n^2 \mod 4$ .

### Division and remainder

If  $n \in \mathbb{Z}$  is odd, then  $n^2 \mod 8 = 1$ . Equivalently,

 $\forall$  odd n,  $\exists m \in \mathbb{Z}$  such that  $n^2 = 8m + 1$ .

### Definition

Given  $x \in \mathbb{R}$ , the floor of x is defined as

 $|x| = n \quad \Leftrightarrow \quad n \leq x < n+1.$ 

The ceiling of x is defined as

$$[x] = n \quad \Leftrightarrow \quad n - 1 < x \le n.$$

### Questions

Are the following true or false?

1. 
$$|x-1| = |x| - 1$$

2. 
$$\lfloor x - y \rfloor = \lfloor x \rfloor - \lfloor y \rfloor$$
.

#### Proposition

For all  $x \in \mathbb{R}$  and  $m \in \mathbb{Z}$ ,  $\lfloor x + m \rfloor = \lfloor x \rfloor + m$ .

#### Proof

By definition,  $n \le x < n+1$ , where  $\lfloor x \rfloor = n$ .

Adding m yields

$$\underbrace{m+m}_{|x+m|} \le x+m < n+m+1.$$

Note that  $\lfloor x \rfloor = n$  implies that  $n + m = \lfloor x \rfloor + m$ .

### Proposition

For all integers n,

$$\left\lfloor \frac{n}{2} \right\rfloor = \begin{cases} \frac{n}{2} & n \text{ is even} \\ \frac{n-1}{2} & n \text{ is odd.} \end{cases}$$

### Proposition

For all integers n and d,

$$n \text{ div } d = \left\lfloor \frac{n}{d} \right\rfloor, \quad \text{and} \quad n \text{ mod } d = n - d \left\lfloor \frac{n}{d} \right\rfloor.$$