

## TOPICS: CAUCHY-EULER EQUATIONS AND POWER SERIES SOLUTIONS TO ODES

1. For each of the *Cauchy-Euler equations* below, look for a solution of the form  $y(x) = x^r$ , and plug this back in and find  $r$ . Find a basis of the solution space consisting of two real-valued functions, and use this to write the general solution.

(a)  $x^2y'' - xy' - 3y = 0$

(b)  $x^2y'' - xy' + 5y = 0$

(c)  $x^2y'' - 3xy' + 4y = 0$

2. Write each of the following as a single series of the form  $\sum f(n)x^n$ . That is,  $f(n)$  is the coefficient of  $x^n$ . You may need to additionally “pull out” the first term(s) from one of the sums.

(a)  $\sum_{n=0}^5 x^{n-1}$

(d)  $\sum_{n=0}^{\infty} a_n x^n + \sum_{n=0}^{\infty} b_n x^{n-1} + \sum_{n=0}^{\infty} c_n x^{n+1}$

(b)  $\sum_{n=0}^5 x^{n+1}$

(e)  $5 \sum_{n=0}^{\infty} n(n-1)x^{n-2} + \sum_{n=0}^{\infty} nx^{n-1} - \sum_{n=0}^{\infty} x^n$ .

(c)  $\sum_{n=0}^{\infty} na_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n$

3. Consider the ODE  $y'' - 2xy' + 10y = 0$ . Note that unlike the equation in the first problem, there will not longer be a simple solution of the form  $x^r$ . However, we know that the solution space is 2-dimensional, and most “nice” functions can be written as a power series. Therefore, we’ll look for a solution of the form  $y(x) = \sum_{n=0}^{\infty} a_n x^n$ .

(a) Plug  $y(x)$  back into the ODE and find a recurrence relation for  $a_{n+2}$  in terms of  $a_n$  and  $a_{n+1}$ .

(b) Explicitly write out the coefficients  $a_n$  for  $n \leq 9$ , in terms of  $a_0$  and  $a_1$ . Write down formulas for  $a_{2n}$  and  $a_{2n+1}$  in terms of  $a_0$  and  $a_1$ .

(c) Since the solution space to this ODE is 2-dimensional, the general solution you found in Part (a) can be written as  $y(x) = C_0y_0(x) + C_1y_1(x)$ . Find such a *basis*,  $\{y_0(x), y_1(x)\}$ .

(d) Find a non-zero *polynomial* solution. [*Hint*: Make a good choice for  $a_0$  and  $a_1$ .]

(e) Are there any other polynomial solutions, excluding scalar multiples of the one you found in (d)? Why or why not?

(f) Consider the initial value problem

$$y'' - 2xy' + 10y = 0, \quad y(0) = x_0, \quad y'(0) = v_0.$$

What are  $x_0$  and  $v_0$  in terms of the coefficients  $a_n$ ?

4. The differential equation  $(1 - x^2)y'' - 2xy' + \nu(\nu + 1)y = 0$ , where  $\nu$  is a constant, is known as *Legendre's equation*. It is used for modeling spherically symmetric potentials in the theory of Newtonian gravitation and in electricity and magnetism.
- Assume that the general solution has the form  $y(t) = \sum_{n=0}^{\infty} a_n x^n$ , and find the recursion formula for  $a_{n+2}$  in terms of  $a_n$  and  $a_{n+1}$ .
  - Use the recursion formula to determine  $a_n$  in terms of  $a_0$  and  $a_1$ , for  $2 \leq n \leq 9$ .
  - For each  $\nu \in \mathbb{N}$ , there will be a single (up to scalar multiples) nonzero polynomial solution  $P_\nu(x)$ , called the *Legendre polynomial* of degree  $\nu$ . Find the Legendre polynomial of degree  $\nu = 3$ .
  - Find a basis for the solution space to  $(1 - x^2)y'' - 2xy' + 12y = 0$ .
5. The differential equation  $y'' - xy = 0$  is called *Airy's equation*, and is used in physics to model the refraction of light.
- Assume a power series solution, and find the recurrence relation of the coefficients. [*Hint*: When shifting the indices, one way is to let  $m = n - 3$ , then factor out  $x^{n+1}$  and find  $a_{n+3}$  in terms of  $a_n$ . Alternatively, you can find  $a_{n+2}$  in terms of  $a_{n-1}$ .]
  - Show that  $a_2 = 0$ . [*Hint*: the two series for  $y''$  and  $xy$  don't "start" at the same power of  $x$ , but for any solution, each term must be zero. (Why?)]
  - Find the particular solution when  $y(0) = 1$ ,  $y'(0) = 0$ , as well as the particular solution when  $y(0) = 0$ ,  $y'(0) = 1$ .

6. Consider the following initial value problem:

$$y''' - y = 0, \quad y(0) = 1, \quad y'(0) = 0, \quad y''(0) = 0.$$

- Assume there is a solution of the form  $y(t) = e^{rt}$ , and plug this back in and solve for  $r$ . Use this to write the general solution. [*Hint*: The equation  $r^3 = 1$  has 3 distinct solutions over  $\mathbb{C}$ , called the *3rd roots of unity*:  $r_1 = e^{0\pi i/3} = 1$ ,  $r_2 = e^{2\pi i/3}$ , and  $r_3 = e^{4\pi i/3}$ .]
- From here, you could solve the initial value problem by plugging in the initial conditions, but you'll get a system of three equations with three unknowns, and involving complex numbers. Here's another way: look for a power series solution,  $y(t) = \sum_{n=0}^{\infty} a_n t^n$ . Plug this back into the ODE and find the recurrence relation for the coefficients.
- Compute  $a_n$  for  $n \leq 10$ , and once you see the pattern, write down the general solution to the ODE as a power series. [*Hint*: It should look familiar!]
- Write down a *basis* for the solution space.
- Plug in the initial conditions and find the particular solution to the IVP.
- Solve a similar initial value problem:

$$y''' - y = 0, \quad y(0) = 1, \quad y'(0) = 1, \quad y''(0) = 1.$$