
Read Chapters 9.2–9.4 of Visual Group Theory (VGT), or Chapters 13–14 of AATA. Then write up solutions to the following exercises.

1. Let S be the following set of 7 "binary squares":



- (a) Consider the (right) action of the group $G = V_4 = \langle v, h \rangle$ on S, where $\phi(v)$ reflects each square vertically, and $\phi(h)$ reflects each square horizontally. Draw an action diagram and compute the stabilizer of each element.
- (b) Consider the (right) action of the group $G = C_4 = \langle r | r^4 = e \rangle$ on S, where $\phi(r)$ rotates each square 90° clockwise. Draw an action diagram and compute the stabilizer of each element.
- (c) Suppose a group G of order 15 acts on S. Show that there must be a fixed point.
- 2. Let $G = S_4$ act on itself by conjugation via the homomorphism

 $\phi \colon G \longrightarrow \operatorname{Perm}(S), \qquad \phi(g) = \operatorname{the permutation that sends each } x \mapsto g^{-1}xg.$

- (a) How many orbits are there? Describe them as specifically as you can.
- (b) Find the orbit and the stabilizer of the following elements:
- i. eiii. (1 2 3)v. (1 2) (3 4)ii. (1 2)iv. (1 2 3 4)
- 3. A *p*-group is a group of order p^k for some integer k. Recall that the *center* of a group G is the set of all elements that commute with everything:

$$Z(G) = \{ z \in G \mid gz = zg, \ \forall g \in G \} = \{ z \in G \mid g^{-1}zg = z, \ \forall g \in G \}.$$

Finally, a group G is *simple* if its only normal subgroups are G and $\langle e \rangle$.

(a) Let G act on itself by conjugation via the homomorphism

 $\phi \colon G \longrightarrow \operatorname{Perm}(S), \qquad \phi(g) = \operatorname{the permutation that sends each } x \mapsto g^{-1}xg.$

Show that $Fix(\phi) = Z(G)$.


- (b) Show that if G is a p-group, then |Z(G)| > 1. [Hint: Revisit the Class Equation.]
- (c) Use the result of the previous part to classify all simple p-groups.

- 4. Let G be an unknown group of order 8. By the First Sylow Theorem, G must contain a subgroup H of order 4.
 - (a) If all subgroups of G of order 4 are isomorphic to V_4 , then what group must G be? Completely justify your answer.
 - (b) Next, suppose that G has a subgroup $H \cong C_4$. Then G has a Cayley diagram like one of the following:



Find all possibilities for finishing the Cayley diagram.

- (c) Label each completed Cayley diagram by isomorphism type. Justify your answer.
- (d) Make a complete list of all groups of order 8, up to isomorphism.
- 5. In this problem, we will find the Sylow subgroups of all 3 nonabelian groups of order 12.
 - (a) Find all Sylow 2-subgroups and Sylow 3-subgroups of the groups whose subgroup lattices are shown below. Determine which are normal



- (b) Find all Sylow subgroups of $D_6 = \langle r, f \mid r^6 = f^2 = e, rfr = f \rangle$, and determine which are normal.
- 6. Recall that a group G is called *simple* if its only normal subgroups are G and $\{e\}$. Use group actions and/or the Sylow theorems to show the following.
 - (a) There is no simple group of order $45 = 3^2 \cdot 5$.
 - (b) There is no simple group of order pq, where p < q and are both prime.
 - (c) There is no simple group of order $56 = 2^3 \cdot 7$.
 - (d) There is no simple group of order $108 = 2^2 \cdot 3^3$.
 - (e) If G has a subgroup H with [G : H] = p, the smallest prime dividing |G|, then $H \leq G$, and hence G cannot be simple. [Hint: Let G act on the right cosets of H.]