

**MTHSC 208 (Differential Equations)**  
**Dr. Matthew Macauley**  
**HW 19**  
**Due Friday April 3rd, 2009**

- (1) The function

$$f(t) = \begin{cases} 0 & -\pi \leq t < \pi/2, \\ 1 & -\pi/2 \leq t < \pi/2, \\ 0 & \pi/2 \leq t \leq \pi, \end{cases}$$

can be extended to be periodic of period  $2\pi$ . Sketch the graph of the resulting function, and compute its Fourier series.

- (2) The function

$$f(t) = |t|, \quad \text{for } t \in [-\pi, \pi]$$

can be extended to be periodic of period  $2\pi$ . Sketch the graph of the resulting function, and compute its Fourier series.

- (3) The function

$$f(t) = \begin{cases} 0 & -\pi \leq t < 0, \\ t & 0 \leq t \leq \pi, \end{cases}$$

can be extended to be periodic of period  $2\pi$ . Sketch the graph of the resulting function, and compute its Fourier series.

- (4) Consider the  $2\pi$ -periodic function defined by

$$f(t) = \begin{cases} t^2 & -\pi \leq t < \pi, \\ f(t - 2k\pi), & -\pi + 2k\pi \leq t < \pi + 2k\pi. \end{cases}$$

Sketch this function and compute its Fourier series.

- (5) Find the Fourier series of the function  $f(t) = 2 - 3\sin 4t + 5\cos 6t$ , and sketch the graph of this function (use your calculator). *Hint: this problem is simple – don't do any integrals!*
- (6) Sketch the graph of the function  $f(t) = \sin^2 t$  and find its Fourier series. *Hint: Don't do any integrals! Instead, use a standard trig identity.*
- (7) Which functions from the previous exercises had only cosine terms in their Fourier series expansion? Which functions only had sine terms? Which had both? Do you see a pattern? *Hint: compare the symmetries of the graphs of these functions to the symmetries of the graphs of sine waves and cosine waves.*