MthSc 208, Fall 2011 (Differential Equations) Dr. Macauley

HW 15

Due Monday April 4th, 2011

- (1) For each initial value problem, sketch the forcing term, and then solve for y(t). Write your solution as a piecewise function (i.e., not using the Heavysie function). Recall that the function $H_{ab}(t) = H(t-a) - H(t-b)$ is the interval function.

 - (a) $y'' + 4y = H_{01}(t)$, y(0) = 0, y'(0) = 0(b) $y'' + 4y = t H_{01}(t)$, y(0) = 0, y'(0) = 0
- (2) Define the function

$$\delta_p^{\epsilon}(t) = \frac{1}{\epsilon} \left(H_p(t) - H_{p+\epsilon}(t) \right) .$$

(a) Show that the Laplace transform of $\delta_p^{\epsilon}(t)$ is given by

$$\mathcal{L}\left\{\delta_p^{\epsilon}(t)\right\} = e^{-sp} \frac{1 - e^{-s\epsilon}}{s\epsilon}.$$

- (b) Use l'Hôpital's rule to take the limit of the result in part (a) as $\epsilon \to 0$. How does this result agree with the fact that $\mathcal{L}\{\delta_p(t)\}=e^{-sp}$?
- (3) Use a Laplace transform to solve the follosing initial value problem:

$$y' = \delta_p(t), \qquad y(0) = 0$$

How does your answer support what engineers like to say, that the "derivative of a unit step is a unit impulse"?

(4) Define the function

$$H_p^{\epsilon}(t) = \left\{ \begin{array}{ll} 0, & 0 \leq t$$

- (a) Sketch the graph of $H_p^{\epsilon}(t)$.
- (b) Without being too precise about things, we could argue that $H_p^{\epsilon}(t) \to H_p(t)$ as $\epsilon \to 0$, where $H_p(t) = H(t-p)$. Sketch the graph of the derivative of $H_p^{\epsilon}(t)$.
- (c) Compare your result in (b) with the graph of $\delta_p^{\epsilon}(t)$. Argue that $H_p'(t) = \delta_p(t)$.
- (5) Solve the following initial value problems.
 - (a) $y'' + 4y = \delta(t)$, y(0) = 0, y'(0) = 0
 - (b) $y'' 4y' 5y = \delta(t)$, y(0) = 0, y'(0) = 0