1. Let  $\mathcal{P}_2$  be the vector space of all polynomials  $p(x) = a_0 + a_1 x + a_2 x^2$  over  $\mathbb{R}$ , with degree  $\leq 2$ . Let  $\xi_1, \xi_2, \xi_3$  be distinct real numbers, and define

$$\ell_j = p(\xi_j) \text{ for } j = 1, 2, 3.$$

- (a) Show that  $\ell_1, \ell_2, \ell_3$  are linearly independent functions on  $\mathcal{P}_2$ .
- (b) Show that  $\ell_1, \ell_2, \ell_3$  is a basis for the dual space  $\mathcal{P}'_2$ .
- (c) Find polynomials  $p_1(x), p_2(x), p_3(x)$  in  $\mathcal{P}_2$  of which  $\ell_1, \ell_2, \ell_3$  is the dual basis in  $\mathcal{P}'_2$ .
- 2. Let W be the subspace of  $\mathbb{R}^4$  spanned by (1, 0, -1, 2) and (2, 3, 1, 1). Which linear functions  $\ell(x) = c_1 x_1 + c_2 x_2 + c_3 x_3 + c_4 x_4$  are in the annihilator of W? Write your answer by giving an explicit basis of  $W^{\perp}$ .
- 3. Let  $T: X \to U$  be a linear map. Prove the following:
  - (a) The image of a subspace of X is a subspace of U.
  - (b) The inverse image of a subspace of U is a subspace of X.
- 4. Prove Theorem 3.3 in Lax:
  - (a) The composite of linear mappings is also a linear mapping.
  - (b) Composition is distributive with respect to the addition of linear maps, that is,


$$(R+S) \circ T = R \circ T + S \circ T$$

and

$$S \circ (T+P) = S \circ T + S \circ P,$$

where  $R, S: U \to V$  and  $P, T: X \to U$ .

- 5. Let X be a finite-dimensional vector space over K and let  $\{x_1, \ldots, x_n\}$  be an ordered basis for X. Let U be a vector space over the same field K but possibly with a different dimension, and let  $\{u_1, \ldots, u_n\}$  be an arbitrary set of vectors in U. Show that there is precisely one linear transformation  $T: X \to U$  such that  $Tx_i = u_i$  for each  $i = 1, \ldots, n$ .
- 6. Let X and U be vector spaces, and suppose that Y is a subspace of X. Let  $Q: X \to X/Y$  be the canonical quotient map sending  $x \xrightarrow{Q} \{x\}$ , and let  $T: X \to U$  be a linear map. Give necessary and sufficient conditions for the existence of a unique linear map  $S: X/Y \to U$  such that  $T = S \circ Q$ . When this happens, the map T is said to factor through the quotient space, as shown by the following commutative diagram:



Prove all of your claims.