

Read: Lax, Chapter 7, pages 89–100.

1. Let X be a finite-dimensional real Euclidean space. We say that a sequence $\{A_n\}$ of linear maps converges to a limit A if $\lim_{n \rightarrow \infty} \|A_n - A\| = 0$.
 - (a) Show that $\{A_n\}$ converges to A if and only if for all $x \in X$, $A_n x$ converges to Ax .
 - (b) Show by example that this fails if $\dim X = \infty$.
2. Let $A: X \rightarrow U$ be a linear map between Euclidean spaces, and let $A^*: U \rightarrow X$ denote the adjoint map. The map A has a *left inverse* if there is a linear map $L: U \rightarrow X$ such that $LA = I_X$, the identity on X . It has a *right inverse* if there is a linear map $R: U \rightarrow X$ such that $AR = I_U$ is the identity on U .
 - (a) Prove that $R_{A^*}^\perp = N_A$.
 - (b) Prove that A maps R_{A^*} bijectively onto R_A .
 - (c) Show that if A has a left inverse, then $Ax = u$ has *at most* one solution. Give a condition on u that completely characterizes when there is a solution.
 - (d) Show that if A has a right inverse, then $Ax = u$ has *at least* one solution. If $Ax_p = u$ for some particular $x_p \in X$, then describe all solutions for x in this case. What condition ensures that there will be *only* one solution?
 - (e) What are the possibilities for the rank of A if it has a left inverse? What if it has a right inverse?
3. Let X be the space of continuous complex-valued functions on $[-1, 1]$ and define an inner product on X by

$$(f, g) = \int_{-1}^1 f(s)\bar{g}(s) ds.$$

Let $m(s)$ be a continuous function of absolute value 1, that is, $|m(s)| = 1$, $-1 \leq s \leq 1$. Define M to be multiplication by m :

$$(Mf)(s) = m(s)f(s).$$

Show that M is unitary.

4. Let A be a linear map of a finite-dimensional complex Euclidean space X .
 - (a) A matrix is *normal* if $AA^* = A^*A$. It is unitarily similar to a diagonal matrix if $A = U^*DU$ for a diagonal matrix D and unitary matrix U . Show that these conditions are equivalent.
 - (b) Prove that if A is normal then it has a square-root, that is, a matrix B such that $A = B^2$. Is B necessarily normal? Unique?
 - (c) Suppose that A is diagonalizable. Prove that A is normal if and only if each eigenvector of A is an eigenvector of A^* .

5. Express $q(x_1, x_2, x_3) = 3x_1^2 + 8x_1x_2 - 7x_1x_3 + 12x_2^2 - 8x_2x_3 + 6x_3^2$ as $q(x) = x^T Ax$, where A is symmetric.

6. Let

$$M = \begin{bmatrix} 3 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 3 \end{bmatrix},$$

and let $q(x) = (x, Mx)$. Find an orthogonal matrix P which diagonalizes the quadratic form q .