

*Read:* Strang, Section 3.1, 3.2.

*Suggested short conceptual exercises:* Strang, Section 3.1, #15, 16, 23, 25–27, 29. Section 3.2, #3, 6, 9, 10–18, 20, 28–31.

1. For each subsets of  $\mathbb{R}^3$ , determine if it is a subspace. If not, give an explicit example of how it fails.
  - (a) The plane of vectors  $\mathbf{b} = (b_1, b_2, b_3)$  with  $b_1 = b_2$ .
  - (b) The plane of vectors with  $b_1 = 1$ .
  - (c) The vectors with  $b_1 b_2 b_3 = 0$ .
  - (d) All linear combinations of  $\mathbf{v} = (1, 4, 0)$  and  $\mathbf{w} = (2, 2, 2)$ .
  - (e) All vectors that satisfy  $b_1 + b_2 + b_3 = 0$ .
  - (f) All vectors with  $b_1 \leq b_2 \leq b_3$ .
2. Let  $P$  be the plane in  $\mathbb{R}^3$  with equation  $x + y - 2z = 4$ , which does not contain the zero-vector, and let  $P_0$  be the parallel plane that passes through the origin.
  - (a) Write the plane  $P_0$  as an equation  $ax + by + cz = d$ .
  - (b) Find two vectors in  $P$  and check that their sum is not in  $P$ .
  - (c) Find two non-colinear vectors in  $P_0$  and check that their sum is in  $P_0$ .
  - (d) Write the set of vectors in  $P_0$  as a linear combination  $c\mathbf{v} + d\mathbf{w}$  for some  $\mathbf{v}$  and  $\mathbf{w}$ .
  - (e) Write the set of vectors in  $P$  as  $\mathbf{x}_p + c\mathbf{v} + d\mathbf{w}$ . That is, find such an  $\mathbf{x}_p$  that works.
  - (f) Find a  $3 \times 2$  matrix  $\mathbf{A}$  whose column space of is the plane  $P_0$ .
3. The matrix  $\mathbf{A} = \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix}$  can be thought of as a “vector” in the space  $\mathcal{M}$  of all  $2 \times 2$  matrices.
  - (a) Write down the zero vector in this space, the vector  $\frac{1}{2}\mathbf{A}$ , and the vector  $-\mathbf{A}$ .
  - (b) What matrices are in the smallest subspace containing  $\mathbf{A}$ ?
  - (c) Describe a subspace of  $\mathcal{M}$  that contains  $\mathbf{B} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  but not  $\mathbf{C} = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix}$ .
  - (d) If a subspace of  $\mathcal{M}$  contains  $\mathbf{B}$  and  $\mathbf{C}$ , must it contain  $\mathbf{I}$ ?
  - (e) Describe a nontrivial subspace of  $\mathcal{M}$  that contains no nonzero diagonal matrices.
4. Fix  $n$ , and let  $\mathcal{M}$  be the space of all  $n \times n$  matrices. For either of the following subsets of  $\mathcal{M}$ , determine whether it is a subspace. If true, give a reason, and if false, show why it fails with an explicit example.
  - (a) The set of invertible matrices.

- (b) The set of singular (non-invertible) matrices.
- (c) The set of symmetric matrices ( $\mathbf{A}^T = \mathbf{A}$ ).
- (d) The set of skew-symmetric matrices ( $\mathbf{A}^T = -\mathbf{A}$ ).
- (e) The set of unsymmetric matrices ( $\mathbf{A}^T \neq \mathbf{A}$ ).
5. For each system, determine which right sides (find a condition on  $b_1$ ,  $b_2$ , and  $b_3$ ) that makes them solvable. Describe the column space geometrically (line, plane,  $\mathbb{R}^3$ , etc.) in each case.

$$\begin{bmatrix} 1 & 4 & 2 \\ 2 & 8 & 4 \\ -1 & -4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} \qquad \begin{bmatrix} 1 & 4 \\ 2 & 9 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

6. Consider the following two matrices

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 2 & 4 & 6 \\ 1 & 2 & 3 & 6 & 9 \\ 0 & 0 & 1 & 2 & 3 \end{bmatrix} \qquad \mathbf{B} = \begin{bmatrix} 2 & 4 & 2 \\ 0 & 4 & 4 \\ 0 & 8 & 8 \end{bmatrix}$$

For each of these matrices, carry out the steps outlined below.

- (a) Reduce the matrix to its ordinary echelon form  $\mathbf{U}$ . Determine which are the free variables and which are the pivot variables. What is its rank?
- (b) Find a *special solution* for each free variable. That is, set the free variable to 1 and the other free variables to 0.
- (c) By combining the special solutions, describe every solution to  $\mathbf{Ax} = \mathbf{0}$  and  $\mathbf{Bx} = \mathbf{0}$ . This is the *nullspace* of the matrix.
- (d) Write down the *nullspace matrix*  $\mathbf{N}$ , whose columns are the special solutions.
- (e) By further row operations on each  $\mathbf{U}$ , find the reduced row echelon form  $\mathbf{R}$ .
- (f) Embedded within the nonzero rows of  $\mathbf{R}$  are two (not necessarily contiguous) submatrices, one an identity matrix and the other a matrix  $\mathbf{F}$ . Identify  $\mathbf{F}$ . Now,  $\mathbf{N}$  consists of two (not necessarily contiguous) submatrices: what are they?
- (g) True or false: The nullspace of  $\mathbf{R}$  equals the nullspace of  $\mathbf{U}$ ?
7. For each of the following part, construct a matrix  $\mathbf{A}$  with the specific properties specified. In this problem,  $n \times 1$  column vectors are written as ordered  $n$ -tuples.
- (a) The column space contains  $(1, 1, 1)$  and the nullspace is the line of multiples of  $(1, 1, 1, 1)$ .
- (b)  $\mathbf{A}$  is a  $2 \times 2$  matrix whose column space equals its nullspace.
- (c) The nullspace of  $\mathbf{A}$  consists of all linear combinations of  $(2, 2, 1, 0)$  and  $(3, 1, 0, 1)$ . [Hint: Try a  $2 \times 4$  matrix, and think of the “column picture.”]
- (d) The nullspace of  $\mathbf{A}$  consists of all multiples of  $(4, 3, 2, 1)$ .
- (e) The column space contains  $(1, 1, 5)$  and  $(0, 3, 1)$  and the nullspace contains  $(1, 1, 2)$ .