

Read: Lax, Chapter 5, pages 44–57.

1. Let  $S_n$  denote the set of all permutations of  $\{1, \dots, n\}$ .
  - (a) Prove that  $\text{sgn}(\pi_1 \circ \pi_2) = \text{sgn}(\pi_1) \text{sgn}(\pi_2)$ .
  - (b) Let  $\pi \in S_n$ , and suppose that  $\pi = \tau_k \circ \dots \circ \tau_1 = \sigma_\ell \circ \dots \circ \sigma_1$ , where  $\tau_i, \sigma_j \in S_n$  are transpositions. Prove that  $k \equiv \ell \pmod{2}$ .
2. Let  $X$  be an  $n$ -dimensional vector space over a field  $K$ .
  - (a) Prove that if the characteristic of  $K$  is not 2, then every skew-symmetric form is alternating.
  - (b) Give an example of a non-alternating skew-symmetric form.
  - (c) Give an example of a non-zero alternating  $k$ -linear form ( $k < n$ ) such that  $f(x_1, \dots, x_k) = 0$  for some set of linearly independent vectors  $x_1, \dots, x_k$ .
3. Let  $X$  be a 2-dimensional vector space over  $\mathbb{C}$ , and let  $f: X \times X \rightarrow \mathbb{C}$  be an alternating, bilinear form. If  $\{x_1, x_2\}$  is a basis of  $X$ , determine a formula for  $f(u, v)$  in terms of  $f(x_1, x_2)$ , and the coefficients used to express  $u$  and  $v$  with this basis. [Pun intended!]
4. Let  $X$  be an  $n$ -dimensional vector space over  $\mathbb{R}$ , and let  $f$  be a non-degenerate symmetric bilinear form. That is, it has the additional property that for all nonzero  $x \in X$ , there is some  $y \in X$  for which  $f(x, y) \neq 0$ .
  - (a) Prove that the map  $L: X \rightarrow X'$  given by  $L: x \mapsto f(x, -)$  is an isomorphism.
  - (b) Show that, given any basis  $x_1, \dots, x_n$  for  $X$ , there exists a basis  $y_1, \dots, y_n$  such that  $f(x_i, y_j) = \delta_{ij}$ .
  - (c) Conversely, prove that if  $\mathcal{B}_X = \{x_1, \dots, x_n\}$  and  $\mathcal{B}_Y = \{y_1, \dots, y_n\}$  are sets of vectors in  $X$  with  $f(x_i, y_j) = \delta_{ij}$ , then  $\mathcal{B}_X$  and  $\mathcal{B}_Y$  are bases for  $X$ .
5. Let  $X$  be an  $n$ -dimensional vector space over  $\mathbb{R}$ , and let  $f$  be a non-degenerate symmetric bilinear form.
  - (a) Show that there exists  $x_1 \in X$  with  $f(x_1, x_1) \neq 0$ .
  - (b) Any fixed  $x_1 \in X$  for which  $f(x_1, x_1) \neq 0$  induces a *linear* map  $T = f(x_1, -)$ . Show that the nullspace  $N_T$  has dimension  $n - 1$ .
  - (c) Let  $Z_1 = N_T$ . Show that the restriction of  $f$  to  $Z_1 \times Z_1$  is again non-degenerate.
  - (d) Prove that  $X$  has a basis  $\{x_1, \dots, x_n\}$  such that  $f(x_i, x_i) \neq 0$  for all  $i$ , and  $f(x_i, x_j) = 0$  whenever  $i \neq j$ .
  - (e) Give an example of a vector space  $X$  ( $2 \leq \dim X < \infty$ ) with basis  $\mathcal{B}$  and a non-degenerate symmetric bilinear form  $f$  for which  $f(x, x) = 0$  for all  $x \in \mathcal{B}$ .
6. Let  $A = (c_1, \dots, c_n)$  be an  $n \times n$  matrix ( $c_i$  is a column vector), and let  $B$  be the matrix obtained from  $A$  by adding  $k$  times the  $i^{\text{th}}$  column of  $A$  to the  $j^{\text{th}}$  column of  $A$ , for  $i \neq j$ . Prove that  $\det A = \det B$ .