

Lecture 3.9: The method of Frobenius

Matthew Macauley

Department of Mathematical Sciences
Clemson University
<http://www.math.clemson.edu/~macaule/>

Math 2080, Differential Equations

Quick review of power series

Definitions

A **power series** centered at x_0 is a limit of **partial sums**:

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n = \lim_{N \rightarrow \infty} \sum_{n=0}^N a_n(x - x_0)^n.$$

It **converges** at x if the sequence of partial sums converges. Otherwise, it **diverges**.

Examples

- The power series $\lim_{N \rightarrow \infty} \sum_{n=0}^N \frac{1}{n!} x^n$ **converges** to e^x for all $x \in (-\infty, \infty)$.
- The power series $\lim_{N \rightarrow \infty} \sum_{n=0}^N (-1)^n x^n$ **converges** to $\frac{1}{1+x}$ for all $x \in (-1, 1)$. It **diverges** at $x = 1$.

Radius of convergence

The largest number R such that if $|x - x_0| < R$, then $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges.

Ordinary vs. singular points of ODEs

Definitions

A function $f(x)$ is **real analytic** at x_0 if $f(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$ for some $R > 0$.

Definition

Consider the ODE $y'' + P(x)y' + Q(x)y = 0$.

- The point x_0 is an **ordinary point** if $P(x)$ and $Q(x)$ are real analytic at x_0 .
- Otherwise x_0 is a **singular point**, which is:
 - **regular** if $(x - x_0)P(x)$ and $(x - x_0)^2Q(x)$ are real analytic.
 - **irregular** otherwise.

Example

Consider the homogeneous ODEs $y'' + x^2y - 4y = 0$.

Regular vs. irregular singular points

More examples

1. $(1 - x^2)y'' - xy' + p^2y = 0.$

2. $x^3y'' + y' + y = 0.$

When does an ODE have a power series solution?

Theorem of Frobenius

Consider an ODE $y'' + P(x)y' + Q(x)y = f(x)$. If x_0 is an **ordinary point**, and $P(x)$, $Q(x)$, and $f(x)$ have radii of convergence R_P , R_Q , and R_f , respectively, then there is a **power series solution**

$$y(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n, \quad R = \min\{R_P, R_Q, R_f\}.$$

If x_0 is a **regular singular point** and $(x - x_0)P(x)$, $(x - x_0)^2Q(x)$, and $f(x)$ have radii of convergence R_P , R_Q , and R_f , respectively, then there is a **generalized power series solution**

$$y(x) = (x - x_0)^r \sum_{n=0}^{\infty} a_n(x - x_0)^n, \quad R = \min\{R_P, R_Q, R_f\},$$

for some constant r .

An ODE with a generalized power series solution

Example 5

Solve the homogeneous differential equation $2xy'' + y' + y = 0$.

Applications of the power series method

Examples from physics and engineering

- Cauchy-Euler equation: $x^2y'' + axy' + by = 0$. Arises when solving Laplace's equation in polar coordinates.
- Hermite's equation: $y'' - 2xy' + 2py = 0$. Used for modeling simple harmonic oscillators in quantum mechanics.
- Legendre's equation: $(1 - x^2)y'' - 2xy' + p(p + 1)y = 0$. Used for modeling spherically symmetric potentials in the theory of Newtonian gravitation and in electricity & magnetism (e.g., the wave equation for an electron in a hydrogen atom).
- Bessel's equation: $x^2y'' + xy' + (x^2 - p^2)y = 0$. Used for analyzing vibrations of a circular drum.
- Airy's equation: $y'' - k^2xy = 0$. Models the refraction of light.
- Chebyshev's equation: $(1 - x^2)y'' - xy' + p^2y = 0$. Arises in numerical analysis techniques.