

## TOPICS: SELF-ADJOINT OPERATORS AND STURM-LIOUVILLE THEORY

For consistency, we will say that a *Sturm-Liouville equation* is a second-order differential equation in the following *self-adjoint form*:

$$-\frac{d}{dx}(p(x)y') + q(x)y = \lambda w(x)y, \quad (1)$$

where  $p(x) > 0$  and  $w(x) > 0$  is called the *weight*, or *density* function. If we divide through by  $w(x)$ , we can write this equation as  $Ly = \lambda y$ , where  $L$  is a *self-adjoint* linear operator. The possible values of  $\lambda$  are the *eigenvalues*, and solutions are the *eigenfunctions*.

1. Write the following differential equations in self-adjoint form. That is, put them in the above form, and find  $p(x)$ ,  $q(x)$ , and the *weight*  $w(x)$ . Also, write out the corresponding linear operator  $L$ .

(a) Airy's equation:  $y'' + (\lambda - x)y = 0$

(b) Laguerre's equation:  $xy'' + (1 - x)y' + \lambda y = 0$

(c) An arbitrary linear equation:  $y'' + P(x)y' + Q(x)y = \lambda R(x)y$ . [*Hint*: Multiply through by an *integrating factor*,  $e^{\int P(x)dx}$ .]

2. In this problem, we will find all solutions to the Sturm-Liouville problem

$$-y'' = \lambda y, \quad y'(0) = y(L) = 0.$$

(a) First, suppose that  $\lambda = 0$ . That is, solve  $y'' = 0$ ,  $y'(0) = y(L) = 0$ .

(b) Next, suppose  $\lambda = -\omega^2 \leq 0$ . That is, solve the boundary value problem  $y'' = \omega^2 y$ ,  $y'(0) = y(L) = 0$ . [*Hint*: When the domain is finite, e.g.,  $[0, L]$ , it is usually more convenient to use cosh and sinh instead of exponentials.]

(c) Finally, suppose  $\lambda = \omega^2 > 0$ . That is, solve  $y'' = -\omega^2 y$ ,  $y'(0) = y(L) = 0$ .

(d) Summarize the results from Parts (a)–(c) in terms of the eigenvalues and corresponding eigenfunctions of a particular linear differential operator  $L$ . What is  $L$ ?

(e) Sketch the first four eigenfunctions on  $[0, L]$ .

3. By the main theorem of Sturm-Liouville theory, if we define an inner product as

$$\langle f, g \rangle = \int_a^b f(x)\overline{g(x)}w(x) dx, \quad (2)$$

then the eigenfunctions  $\{y_n(x)\}$  form an *orthogonal basis* (Note: not necessarily *orthonormal*!) for the space of functions, integrable on  $[a, b]$  with  $\langle f, f \rangle < \infty$  that satisfy the boundary conditions. This means that for any  $f \in L^2([a, b], w)$  with the same boundary conditions, we can write

$$f(x) = \sum_{n=1}^{\infty} c_n y_n(x).$$

- (a) Consider the Sturm-Liouville problem from Part (c) of the previous problem:

$$-y'' = \lambda y, \quad y'(0) = 0, \quad y(L) = 0.$$

What is  $w(x)$ ?

- (b) The function  $f(x) = x^2 - L^2$  is clearly continuous and satisfies  $f'(0) = f(L) = 0$ . Compute the *norm*  $\|f\| := \langle f, f \rangle^{1/2}$  of  $f$ .
- (c) Since the eigenfunctions form a basis for the subspace of  $L^2([0, L]; w)$  that satisfy the above boundary conditions, we can write

$$x^2 - L^2 = \sum_{n=1}^{\infty} c_n y_n(x), \quad 0 \leq x \leq L.$$

Write down a formula for the  $c_n$ 's. Leave your answer in terms of an integral – no need to actually compute it! [*Hint*: Don't forget that  $y_n(x)$  isn't necessarily of unit length!]

4. Consider the following Sturm-Liouville problem:

$$-y'' - y' = \lambda y, \quad y(0) = 0 \quad y(2) = 0.$$

- (a) Find the eigenvalues and eigenfunctions. [*Hint*: You will encounter a *discriminant* of  $D = 1 - 4\lambda$ . As before, there will be three cases:  $D = 0$ ,  $D > 0$ , and  $D < 0$ .]
- (b) Write this differential equation in standard form, as in Eq. (1). [*Hint*: First, multiply through by an *integrating factor*,  $e^x$ .]
- (c) Write a formula for  $\langle y_n, y_m \rangle$  in terms of an integral. What is this integral equal to when  $n \neq m$ ?
5. Consider the following Sturm-Liouville equation on  $[-1, 1]$ , called *Legendre's differential equation*:

$$(1 - x^2)y'' - 2xy' + \lambda y = 0 \tag{3}$$

In this problem, you will find the eigenvalue and eigenfunctions, which have already come up several times in this class in different settings.

- (a) Write Legendre's equation into *self-adjoint form*, as in Equation 1. That is, find  $p(x)$ ,  $q(x)$ , and  $w(x)$ , and the self-adjoint operator  $L$ . This is called a *singular Sturm-Liouville problem* on the interval  $[a, b] = [-1, 1]$  because the function  $p(x)$  satisfies  $p(-1) = p(1) = 0$ , and so boundary conditions on  $y(x)$  are not needed.
- (b) Assume that there is a power series solution of the form  $\sum_{n=0}^{\infty} a_n x^n$ . Plug this back into Eq. (3) and find the recurrence relation for the coefficients.
- (c) Recall from HW 5 that a generalized power series solution will have radius of convergence  $R = 1$ , i.e., it will be defined on the open interval  $(-1, 1)$ , but *not* on its endpoints,  $a = -1$  or  $b = 1$ . However, if we have a *polynomial* solution (that is, only finitely many non-zero terms, which happens when  $a_{n+2} = 0$  for some  $n$ ), then this will certainly be defined on all of  $[-1, 1]$ . What values of  $\lambda$  lead to a polynomial solution? (These are the *eigenvalues* of  $L$ .)

- (d) The *eigenfunction* for eigenvalue  $\lambda_k$  is a polynomial  $P_k(x)$  called the *Legendre polynomial* of degree  $k$ . (These arose on HW 2 and HW 5.) By Sturm-Liouville theory, they form an *orthogonal basis* of  $L^2([-1, 1])$ , meaning that

$$\langle P_n, P_m \rangle := \int_{-1}^1 P_n(x)P_m(x) dx = 0, \quad n \neq m.$$

Use the recurrence relation to write out the first five Legendre polynomials,  $P_k(x)$ , for  $k = 0, \dots, 4$ . Normalize each one so they form an *orthonormal* set.

- (e) Write the polynomial  $f(x) = 3x^3 - 2x^2 + 4$  using the first four Legendre polynomials. That is, find  $C_0, C_1, C_2$ , and  $C_3$  such that

$$3x^3 - 2x^2 + 4 = C_0P_0(x) + C_1P_1(x) + C_2P_2(x) + C_3P_3(x).$$

*Hint:* This is *very* similar to a problem you did on HW 2!