

# Difference Equations

Matthew Macauley

Department of Mathematical Sciences  
Clemson University  
<http://www.math.clemson.edu/~macaule/>

Math 4500, Spring 2022

## Motivation: Population dynamics

Consider a population of insects that reproduces daily, of size  $P(t)$ :

- *birth rate* is  $f \in [0, \infty)$ ,
- *death rate* is  $d \in [0, 1]$ .

This can be modeled by a simple equation:

$$\Delta P = fP - dP = (f - d)P.$$

Suppose time is *discretized*, e.g., it only takes integer values:  $t = 0, 1, 2, \dots$

Let  $P_t = P(t)$  = population at time  $t$ .

Then  $\Delta P = P_{t+1} - P_t$ , from which it follows that

$$P_{t+1} = P_t + \Delta P = P_t + (f - d)P_t = (1 + f - d)P_t.$$

Letting  $\lambda = 1 + f - d$  (the "*finite growth rate*"), we can write this as  $P_{t+1} = \lambda P_t$ .

## An example

Consider a population of insects that reproduces daily, with the following parameters:

- *initial population*  $P_0 = 300$ ,
- *birth rate*  $f = .03$ ,
- *death rate*  $d = .01$ .

Then the finite growth rate is  $\lambda = 1 + f - d = 1.02$ , and

$$P_1 = (1.02)P_0$$

$$P_2 = (1.02)P_1 = (1.02)^2 P_0$$

$$P_3 = (1.02)P_2 = (1.02)^3 P_0$$

⋮

It is not difficult to see the closed-form solution  $P_t = \lambda^t P_0$ . This is called *exponential growth*.

## What is a difference equation?

### Definition

Let  $Q$  be a quantity defined for all  $t \in \mathbb{N}$ , such that  $Q_{t+1} = F(Q_t)$ , for some function  $F$ .

In the previous example:  $F(x) = \lambda x$ . This is called the *Malthusian model*. It is a *linear* difference equation because  $F(x)$  is linear.

Let's compare difference equations to differential equations:

- Difference equations are *discrete time, continuous space*.
- Differential equations are *continuous time, continuous space*.

### Exercise

Can you think of a model that is discrete time and discrete space? Or continuous time and discrete space?

## Which type of model to use?

### Broad goals

- Find an appropriate model.
- Analyze models that naturally arise.

For example, consider the following three problems to be modeled:

1. Let  $P$  be a population of  $P_0 = 300$  insects with birth rate  $f = .03$  and death rate  $d = .01$ .
2. Let  $P$  be the value of an initial investment of  $P_0 = 300$  dollars with fixed 2% interest rate, i.e.,  $\lambda = 1.02$ .
3. Let  $P$  be a mass of a population of bacteria that is initially  $P_0 = 300$  grams, with growth rate insects with finite growth rate  $\lambda = 1.02$ .

### Exercise

Which of these are more suited for difference equations, and which for differential equations?

## Logistic equation for population growth

Realistically, a population's growth rate isn't constant – it depends on size. (“*density dependent*”).

### Big idea


Analyze  $\Delta P/P =$  per capita growth rate.

- $P$  small:  $\frac{\Delta P}{P}$  large.
- $P$  large:  $\frac{\Delta P}{P}$  small.
- $P$  too large:  $\frac{\Delta P}{P} < 0$ .

Assumptions:

- Let  $r$  be the growth rate when  $P = 0$ . [Technically,  $r = \lim_{P \rightarrow 0^+} \frac{\Delta P}{P}$ .] This is called the *finite intrinsic growth rate*.
- Let  $M$  be the population for which  $\frac{\Delta P}{P} = 0$ . This is called the *carrying capacity*.
- Suppose the growth rate decreases *linearly* with  $P$ .

## Logistic equation for population growth



Since the growth rate decreases *linearly* with  $P$ , basic algebra gives

$$\frac{\Delta P}{P} = -\frac{r}{M}P + r = r \left( 1 - \frac{P}{M} \right).$$

## Logistic equation for population growth

Substituting  $\Delta P = P_{t+1} - P_t$  into  $\frac{\Delta P}{P} = r \left(1 - \frac{P}{M}\right)$ , followed by easy algebra yields the **discrete logistic model**:

$$P_{t+1} = P_t \left(1 + r \left(1 - \frac{P_t}{M}\right)\right).$$

### Model validation

To see if this model is reasonable, the first thing to check are some simple cases:

- $P \ll M \implies 1 - \frac{P}{M} \approx 1 \implies P_{t+1} \approx (1 + r)P_t$ . [Exponential growth!]
- $P \approx M \implies 1 - \frac{P}{M} \approx 0 \implies P_{t+1} \approx P_t$ .

### Exercise

What is  $F(x)$  in the discrete logistic model? [It must satisfy  $P_{t+1} = F(P_t)$ .]




## Solutions of difference equations

Difference equations, though simple, often have *no closed form solution* for  $P_t$ .

However, we can plot the solutions for various initial values  $P_0$ .


Here are some solutions to the equation  $P_{t+1} = P_t + .2P_t(1 - \frac{P_t}{10})$ .



## Cobwebbing


Consider the difference equation  $\Delta P = 0.8P_t \left(1 - \frac{P_t}{10}\right)$ . Or equivalently,  
 $P_{t+1} = F(P_t) = P_t + 0.8P_t \left(1 - \frac{P_t}{10}\right)$ .

We can numerically find  $P_0, P_1, P_2, \dots$  by plotting  $F(x) = x + 0.8x\left(1 - \frac{x}{10}\right)$  and  $y = x$  on the same axes, and then by “cobwebbing”:



## Cobwebbing

Consider another difference equation:  $\Delta P = 1.8P_t \left(1 - \frac{P_t}{10}\right)$ . Or equivalently,  
 $P_{t+1} = F(P_t) = P_t + 1.8P_t \left(1 - \frac{P_t}{10}\right)$ .



## Questions

1. Sketch a plot of several solution curves  $P(t)$  for the difference equations in the previous two examples.
2. What does the spiraling behavior of this cobweb imply about the population  $P(t)$ ?
3. How does this relate to mass-spring systems? [*Hint*: Think about damping.]
4. What features about a population are highlighted in the logistic equation using difference equations that do not arise using differential equations?